

Appendix V5-7D

Doris North Gold Mine Project:
2011 Aquatics Effects Monitoring Program (AEMP)
Marine Expansion Baseline Report



Hope Bay Mining Limited



DORIS NORTH GOLD MINE PROJECT 2011 Aquatic Effects Monitoring Program (AEMP) Marine Expansion Baseline Report



DORIS NORTH GOLD MINE PROJECT 2011 AQUATIC EFFECTS MONITORING PROGRAM (AEMP) MARINE EXPANSION BASELINE REPORT

December 2011
Project #1009-007-05

Citation:

Rescan. 2011. *Doris North Gold Mine Project: 2011 Aquatic Effects Monitoring Program (AEMP) Marine Expansion Baseline Report*. Prepared for Hope Bay Mining Limited by Rescan Environmental Services Ltd.

Prepared for:



Hope Bay Mining Limited

Prepared by:



Rescan™ Environmental Services Ltd.
Vancouver, British Columbia

DORIS NORTH GOLD MINE PROJECT
2011 Aquatic Effects Monitoring Program (AEMP)
Marine Expansion Baseline Report

Executive Summary

Executive Summary

The 2011 Aquatic Effects Monitoring Program (AEMP) Marine Expansion Baseline work was conducted by Rescan Environmental Services Ltd. (Rescan) on behalf of Hope Bay Mining Ltd. (HBML) for the Doris North Project. The Doris North Property is located approximately 125 km southwest of Cambridge Bay, Nunavut, on the south shore of Melville Sound. The nearest communities are Omingmaktok (Bay Chimo; 75 km to the southwest of the property), Cambridge Bay, and Kingaok (Bathurst Inlet; 160 km to the southwest of the property).

This report presents the results from the 2011 AEMP Marine Expansion Baseline program. The primary objective of the 2011 program was to collect marine baseline data relevant to the proposed discharge of treated water from the Tailings Impoundment Area (TIA) into Roberts Bay. Data from this report was used to provide information to support the preparation of the Roberts Bay environmental assessment report that was prepared for the Doris North amendment package (HBML 2011). Data collected in 2011 will be used in the future expanded AEMP should the proposed Doris North amendments be approved.

Currently, the TIA is permitted to be discharged into Doris Creek under the approved Type A Water Licence (2AM-DOH0713). If the proposed modifications to the Doris North Project are approved, additional underground mining at Doris North will encounter deep saline groundwater and saline talik water. HBML is proposing to discharge this additional mine water into the TIA. This saline TIA water could be detrimental to aquatic life if discharged to Doris Creek. A more environmentally appropriate receiving environment is the marine (ocean) environment as the salt content will closely match that of seawater. To accommodate the additional saline TIA water, HBML is proposing to discharge treated TIA water to the marine environment in Roberts Bay via a subsea pipeline and diffuser (HBML 2011).

The 2011 AEMP Marine Expansion Baseline program involved collecting information for the following: the vertical structure and dissolved oxygen content of the water column, water and sediment quality, phytoplankton, and benthic invertebrates. Two distinct marine basins near the Project area were surveyed four times between April and October 2011: Roberts Bay and Reference Bay. Two potential exposure sites were surveyed in Roberts Bay: one at the proposed discharge site (near-field) and one 2 km downstream towards the mouth of Roberts Bay (far-field). One reference site was surveyed in Reference Bay.

The following text provides a brief summary of the various components sampled as part of the 2011 program.

Water Column Structure and Dissolved Oxygen

The water columns in Roberts Bay and Reference Bay were consistently stratified between April and late September 2011, with lower density water sitting above higher density water (i.e., they were two-layer systems). This stratification was weaker during the ice-covered season (April) when there were no freshwater inputs and in late September (open-water) due to low freshwater inputs, decreasing temperatures, and increased wind-driven mixing. This resulted in deep surface mixed layers of 20 to 25 m in September and 30 m in April.

In July and August, the vertical stratification intensified in Roberts and Reference bays due to melting sea ice and increased riverine inputs. The influx of freshwater resulted in shallower pycnoclines (range: 3-10 m) with warmer, fresher water sitting above colder, saltier water at depth. The surface temperatures were warmest in Roberts Bay in August, reaching 10°C.

Both bays had surface layers that were well oxygenated, with surface dissolved oxygen concentrations ranging from about 9 mg/L in August to 12 mg/L in April. Dissolved oxygen concentrations in the bottom waters of Roberts Bay were higher than the minimum concentration of 8.0 mg/L recommended by the Canadian Council of the Ministers of the Environment (CCME) for the protection of marine and estuarine aquatic life. In August, dissolved oxygen concentrations at depths greater than 25 m in Reference Bay were below the guideline of 8.0 mg/L, reaching 6.63 mg/L at 41 m depth.

Light penetration was high during the summer and fall in both Roberts and Reference bays. The euphotic depth (1% light level) was generally deeper than 16 m and usually extended far below the pycnocline. This suggests that net photosynthesis was possible throughout the upper water column in both bays.

Water Quality

Overall, the water quality parameters were similar between Roberts Bay and Reference Bay. Water clarity was high in both bays as total suspended solids were often below analytical detection and turbidity was consistently low. Some nutrients showed vertical concentration gradients, which were most dramatic during July and August. Nitrate and total phosphorus concentrations were generally lower at the surface than in deep waters. During summer, nitrate was usually below analytical detection (>0.006 mg N/L) in the surface layer, while total phosphorus was detectable throughout the water column. Thus, phytoplankton growth was likely controlled by nitrogen availability when sufficient light was present, which is typical of northern, estuarine systems. Vertical gradients of total organic carbon (TOC) exhibited the opposite trend as nutrients as the summer surface concentrations were usually slightly higher than at depth. Vertical gradients characterized by lower nutrients and higher TOC at the surface than at depth are indicative of phytoplankton growth and biological nutrient uptake.

Roberts Bay and Reference Bay can be considered low-metal environments, as all water quality parameters were below their respective CCME guideline for the protection of marine aquatic life. Several metals (e.g., total lead and total zinc) were usually below their analytical detection limit.

Total cyanide was commonly observed at low concentrations during the surveys. The highest concentrations were found in July in the surface waters of Roberts (0.0021 mg/L) and Reference (0.0019 mg/L) bays, and detectable levels were found throughout the Roberts Bay water column in April. Free cyanide made up a very small proportion of the total cyanide pool as there was little detectable free cyanide in either bay.

Sediment Quality

The deep water sediments (42 m) were composed mainly of silt (mean: 54.2%) and clay (mean: 41.3%) in both bays. Roberts Bay sediments had slightly higher mean concentrations of TOC, total chromium, total copper, total lead, and total zinc than the Reference Bay sediments. Conversely, Reference Bay sediments had higher concentrations of total arsenic. The mean arsenic, chromium, and copper levels found in Roberts Bay sediments were higher than CCME interim marine sediment quality guidelines (ISQGs) for these metals. Mean copper levels in Reference Bay were also higher than the ISQG, and the total arsenic concentrations in this bay were particularly high, exceeding both the ISQG and the probable effects level (PEL) for the protection of marine aquatic life.

Phytoplankton

Mean phytoplankton biomass (as chlorophyll *a*) in the Project area was very low at all sites (0.14 to 0.62 $\mu\text{g chl } a/\text{L}$), reaching a maximum of 1.5 $\mu\text{g chl } a/\text{L}$ in Reference Bay in August. The low concentrations reflected the low under-ice light levels in April, and the low surface layer nutrients and stable water column in July and August. In Roberts Bay, the mean phytoplankton biomass was highest

in September ($0.46 \mu\text{g chl } a/\text{L}$), likely due to the wind-driven transport of nutrients into the surface layer during this time.

Benthos

Mean benthos density was slightly higher in the deep Reference Bay sediments ($1,520 \text{ organisms}/\text{m}^2$) than in Roberts Bay ($1,060 \text{ organisms}/\text{m}^2$). Diverse benthos communities were present in both bays, with fewer taxonomic groups being more evenly distributed in Roberts Bay than in Reference Bay. The benthos community in Roberts Bay was largely dominated by families of polychaete worms and small crustaceans, while polychaetes dominated the Reference Bay benthos community.

Acknowledgements

Acknowledgements

Fieldwork was conducted with the enthusiastic and competent assistance of numerous Hope Bay Mining Limited (HBML) field assistants including: Noah Aklah, Robert Maksagak, Leonard Wingnak, and Pamela Gross.

Extensive support was provided by the HBML Environment and Social Responsibility (ESR) Department, the Health, Safety and Loss Prevention (HSLP) Department, and camp management, as well as Great Slave Helicopters, Braden Burry Expediting, and Nuna Logistics.

This report was prepared for HBML by Rescan Environmental Services Ltd. The 2011 AEMP Marine Expansion Baseline fieldwork was conducted by Rescan scientists Soleil Switzer, (M.Sc.), Mike Henry (Ph.D.), Kerry Marchinko (Ph.D., M.Sc.), Craig Hatt (Dip. Eng. Tech.), and Katsky Venter (M.Sc.). The report was written by Mike Henry and Carol Adly (M.Sc.) and reviewed by Deborah Muggli (Ph.D., M.Sc., R.P.Bio.). The project was managed by Deborah Muggli.

DORIS NORTH GOLD MINE PROJECT
2011 Aquatic Effects Monitoring Program (AEMP)
Marine Expansion Baseline Report

Table of Contents

DORIS NORTH GOLD MINE PROJECT 2011 AQUATIC EFFECTS MONITORING PROGRAM (AEMP) MARINE EXPANSION BASELINE REPORT

Table of Contents

Executive Summary	i
Acknowledgements.....	v
Table of Contents	vii
List of Figures	viii
List of Tables	ix
List of Appendices	ix
Glossary and Abbreviations	xi
1. Introduction	1-1
2. Methods	2-1
2.1 Study Area and Sampling Locations	2-1
2.2 2011 Sampling Schedule	2-1
2.3 Sampling Methods	2-2
2.3.1 Physical Oceanography and Light Penetration	2-2
2.3.2 Water Quality	2-6
2.3.3 Sediment Quality	2-8
2.3.4 Phytoplankton.....	2-9
2.3.5 Benthos.....	2-9
2.3.6 Quality Assurance and Quality Control (QA/QC)	2-11
2.3.6.1 Water Quality QA/QC	2-11
2.3.6.2 Sediment Quality QA/QC	2-11
2.3.6.3 Aquatic Biology QA/QC	2-12
3. Results.....	3-1
3.1 Physical Oceanography and Light Penetration	3-1
3.2 Water Quality	3-5
3.3 Sediment Quality	3-5
3.4 Phytoplankton Biomass	3-20
3.5 Benthic Invertebrates.....	3-20

References.....	R-1
-----------------	-----

List of Figures

FIGURE	PAGE
Figure 1-1. Doris North Project Location	1-2
Figure 2.1-1. Marine AEMP Expansion Sampling Locations, Doris North Project, 2011	2-3
Figure 3.1-1. Temperature, Salinity, and Dissolved Oxygen Profiles at Marine AEMP Expansion Site RB1, 2011	3-2
Figure 3.1-2. Temperature, Salinity, and Dissolved Oxygen Profiles at Marine AEMP Expansion Site RB2, 2011	3-3
Figure 3.1-3. Temperature, Salinity, and Dissolved Oxygen Profiles at Marine AEMP Expansion Site REF-Marine 2, 2011	3-4
Figure 3.2-1. pH and Total Alkalinity at Marine AEMP Expansion Sites, 2011	3-6
Figure 3.2-2. Hardness and Total Suspended Solids at Marine AEMP Expansion Sites, 2011	3-7
Figure 3.2-3. Total Ammonia and Nitrate Concentrations at Marine AEMP Expansion Sites, 2011	3-8
Figure 3.2-4. Total Cyanide and Free Cyanide Concentrations at Marine AEMP Expansion Sites, 2011	3-9
Figure 3.2-5. Radium-226 and Total Aluminum Concentrations at Marine AEMP Expansion Sites, 2011	3-10
Figure 3.2-6. Total Arsenic and Total Cadmium Concentrations at Marine AEMP Expansion Sites, 2011	3-11
Figure 3.2-7. Total Copper and Total Iron Concentrations at Marine AEMP Expansion Sites, 2011	3-12
Figure 3.2-8. Total Lead and Total Molybdenum Concentrations at Marine AEMP Expansion Sites, 2011	3-13
Figure 3.2-9. Total Mercury and Methylmercury Concentrations at Marine AEMP Expansion Sites, 2011	3-14
Figure 3.2-10. Total Nickel and Total Zinc Concentrations at Marine AEMP Expansion Sites, 2011 ...	3-15
Figure 3.3-1. Grain Size and Total Organic Carbon Concentrations in Sediments at Marine AEMP Expansion Sites, 2011	3-16
Figure 3.3-2. Total Arsenic, Total Cadmium, Total Chromium, and Total Copper Concentrations in Sediments at Marine AEMP Expansion Sites, 2011	3-17
Figure 3.3-3. Total Mercury, Methylmercury, Total Lead, and Total Zinc Concentrations in Sediments at Marine AEMP Expansion Sites, 2011	3-18
Figure 3.4-1. Phytoplankton Biomass (as Chlorophyll <i>a</i>) at Marine AEMP Expansion Sites, 2011	3-21

Figure 3.5-1. Benthos Density and Taxonomic Composition at Marine AEMP Expansion Sites, 2011...	3-22
Figure 3.5-2. Benthos Richness, Evenness, and Diversity at Marine AEMP Expansion Sites, 2011	3-23
Figure 3.5-3. Benthos Bray-Curtis Index for Marine AEMP Expansion Sites, 2011	3-24

List of Tables

TABLE	PAGE
Table 2.1-1. Marine AEMP Expansion Baseline Sampling Locations, Descriptions, and Purpose, Doris North Project, 2011	2-1
Table 2.2-1. Environmental Components and Sampling Time, AEMP Marine Baseline Expansion, 2011	2-2
Table 2.3-1. Summary of the Marine AEMP Expansion Baseline Sampling Program, Doris North Project, 2011	2-5
Table 2.3-2. 2011 Physical Oceanography Sampling Dates, Marine AEMP Expansion Program	2-5
Table 2.3-3. 2011 Water Quality Sampling Dates and Depths, Marine AEMP Expansion Program	2-6
Table 2.3-4. Marine AEMP Expansion Water Quality Parameters and Realized Detection Limits, Doris North Project, 2011	2-7
Table 2.3-5. 2011 Sediment Quality and Benthic Invertebrate Sampling Dates and Depths, Marine AEMP Expansion Program	2-8
Table 2.3-6. Marine AEMP Expansion Sediment Quality Parameters and Realized Detection Limits, Doris North Project, August 2011	2-8
Table 2.3-7. 2011 Phytoplankton Biomass Sampling Dates and Depths, Marine AEMP Expansion Program	2-10
Table 3.1-1. Marine AEMP Expansion Secchi Depths, Euphotic Zone Depths, and Estimated Pycnocline Depths, Doris North Project, 2011	3-1
Table 3.3-1. Sediment Quality at Marine AEMP Expansion Sites, Percent of Samples in which Concentrations are Higher than CCME Guidelines, 2011	3-19
Table 3.3-2. Sediment Quality at Marine AEMP Expansion Sites, Factor by which Average Concentrations are Higher than CCME Guidelines, 2011	3-19

List of Appendices

Appendix 3.1-1. Temperature and Salinity Data for the Marine AEMP Expansion Sites, Doris North Project, 2011
Appendix 3.1-2. Dissolved Oxygen Concentrations and Percent Saturation for the Marine AEMP Expansion Sites, Doris North Project, 2011
Appendix 3.2-1. Water Quality Data for Marine AEMP Expansion Sites, Doris North Project, 2011

Appendix 3.2-2. QA/QC Blank Data for Marine AEMP Expansion Water Quality Sampling, Doris North Project, 2011

Appendix 3.3-1. Sediment Quality Data for Marine AEMP Expansion Sites, Doris North Project, 2011

Appendix 3.4-1. Phytoplankton Biomass (as Chlorophyll *a*) Data for Marine AEMP Expansion Sites, Doris North Project, 2011

Appendix 3.5-1. Benthos Taxonomy Data for Marine AEMP Expansion Sites, Doris North Project, 2011

Glossary and Abbreviations

Glossary and Abbreviations

Terminology used in this document is defined where it is first used. The following list will assist readers who may choose to review only portions of the document.

Abbreviation/Acronym	Definition
AEMP	Aquatic Effects Monitoring Program
BC	Bray-Curtis Dissimilarity Index (also known as Bray-Curtis Similarity Index) or British Columbia
Benthos	benthic invertebrates
CCME	Canadian Council of Ministers of the Environment
Chl <i>a</i>	chlorophyll <i>a</i>
Chlorophyll <i>a</i>	An essential light harvesting pigment for photosynthetic organisms including phytoplankton. Because of the difficulty in direct measurement of plant carbon, chlorophyll <i>a</i> has become an established routine ‘proxy’ estimate for plant biomass in aquatic studies.
CTD	conductivity, temperature, depth probe
D	Simpson’s Diversity Index
DO	dissolved oxygen
D _s	Secchi depth
E	Simpson’s Evenness Index
EC	Environment Canada
EEM	Environmental Effects Monitoring
ESR	Environment and Social Responsibility Department
F	family richness
HBML	Hope Bay Mining Limited
HSLP	Health, Safety and Loss Prevention Department
ISQG	interim sediment quality guideline
MMER	Metal Mining Effluent Regulations
n/c	not collected
NIRB	Nunavut Impact Review Board
NTU	Nephelometric Turbidity Units
NWB	Nunavut Water Board
PEL	probable effects level

Pycnocline	The structure of a water column is defined by its density profile, with lower density water sitting above higher density water. The depth zone where the density changes most sharply is referred to as the pycnocline.
QA/QC	quality assurance/quality control
Rescan	Rescan Environmental Services Ltd
Salinity	No units, dimensionless. Historically, many units have been assigned to salinity, for example, parts per thousand (ppt or ‰), Practical Salinity Units (PSU), and Practical Salinity Scale (PSS 78). Salinity is defined on the Practical Salinity Scale (PSS) as the conductivity ratio of a sea water sample to a standard KCl solution. As PSS is a ratio, it has no units.
Talik	A layer of year-round unfrozen ground in an area of permafrost where temperatures are above freezing, allowing water to remain in liquid form.
TIA	Tailings Impoundment Area
TOC	total organic carbon
TSS	total suspended solids

DORIS NORTH GOLD MINE PROJECT
2011 Aquatic Effects Monitoring Program (AEMP)
Marine Expansion Baseline Report

1. Introduction

1. Introduction

The Doris North Gold Mine Project (the Project) is located approximately 125 km southwest of Cambridge Bay, Nunavut, on the south shore of Melville Sound. The nearest communities are Omingmaktok (75 km to the southwest of the property), Cambridge Bay, and Kingaok (Bathurst Inlet; 160 km to the southwest of the property). Figure 1-1 provides a general location map for the Doris North Project.

The Nunavut Impact Review Board (NIRB) issued a Project Certificate for the Doris North Gold Project on September 15, 2006 and a Type A Water Licence (Water Licence # 2AM-DOH0713) was issued by the Nunavut Water Board (NWB) on September 19, 2007. A requirement of the Water Licence was to develop and conduct an Aquatic Effects Monitoring Program (AEMP). The AEMP Plan was reviewed by Environment Canada (EC) and approved by the Nunavut Water Board (NWB) in early 2010 (Rescan 2010). The first year of the program was initiated in 2010 (Rescan 2011), and a second year was completed in 2011. This plan followed Environmental Effects Monitoring (EEM) protocols as per the *Metal Mining Effluent Regulations* (MMER 2002).

The current Water Licence permits the discharge of water from the Tailings Impoundment Area (TIA) into Doris Creek, upstream of the waterfall. If the proposed modifications to the Doris North Project are approved (HBML 2011), additional underground mining at Doris North will encounter deep saline groundwater and saline talik water. HBML is proposing to discharge this additional mine water into the TIA (HBML 2011). This saline TIA water could be detrimental to aquatic life if discharged to Doris Creek. A more environmentally appropriate receiving environment is the marine (ocean) environment as the salt content will closely match that of seawater. To accommodate the additional saline TIA water, HBML is proposing to discharge treated TIA water to the marine environment in Roberts Bay via a subsea pipeline and diffuser (HBML 2011). This proposal was part of the Doris North amendment package submitted to NIRB and NWB in November 2011.

The current AEMP will need to be modified if the Doris North amendment package is approved. A proposed expanded program was included in the Doris North amendment package, and a final expanded program will be developed in consultation with EC.

The purpose of collecting AEMP-compatible data in Roberts Bay in 2011 was to provide information to support the preparation of the Roberts Bay environmental assessment report that was prepared for the Doris North amendment package. Data collected in 2011 will also hopefully be used in the future expanded AEMP should the proposed Doris North amendments be approved.

The 2011 AEMP Marine Expansion Baseline program was run concurrently with the approved Doris North AEMP, and followed the same methods (Rescan 2010, 2011). Baseline data were collected for the following parameters as part of the 2011 AEMP Marine Expansion program:

- o water quality and water column structure;
- o sediment quality;
- o primary producers (phytoplankton); and
- o benthic invertebrate community (abundance and taxonomy).



Figure 1-1

An extensive physical oceanographic program was also conducted in Roberts Bay in 2011 to collect ocean current data so the circulation and flushing rate of the inlet could be modelled. Details of this work will be provided in separate Roberts Bay physical oceanography and Roberts Bay circulation reports.

Chapter 2 of this report presents the methods used during the 2011 AEMP Marine Expansion Baseline program, and Chapter 3 presents graphical results. All raw data are provided in appendices. Much of the information collected was used for the Doris North amendment package, and only brief summaries of results are provided in this report.

2. Methods

2. Methods

2.1 STUDY AREA AND SAMPLING LOCATIONS

The AEMP Marine Expansion Baseline study area included areas in Roberts Bay anticipated to be potentially influenced by the treated TIA discharge and an area beyond any TIA discharge influence (reference area). Two potential exposure sites (RB1 and RB2) in Roberts Bay and one reference site (REF-Marine 2) were sampled in 2011 (Figure 2.1-1). The two exposure sites were situated to acquire baseline information downstream of the proposed treated TIA discharge location. RB1 was considered a near-field exposure site and was in the immediate proposed receiving area (~100 m) of the proposed diffuser. RB2 was considered a far-field exposure site and was 2 km north of the proposed discharge location and 2.3 km from the mouth of Roberts Bay to Melville Sound. The reference site REF-Marine 2 was located in southern Reference Bay for comparability with the two exposure sites in Roberts Bay. This reference site was sufficiently far away from the influence of mining activity and resembled, as much as possible, the hydrological and habitat features of the exposure areas. Figure 2.1-1 shows the sampling sites, the parameters collected, and the Doris North Project infrastructure as of 2011. It also shows the location of the proposed TIA discharge pipeline. A summary table (Table 2.1-1) provides details for the rational of the site selections.

Table 2.1-1. Marine AEMP Expansion Baseline Sampling Locations, Descriptions, and Purpose, Doris North Project, 2011

AEMP Expansion Site	Coordinates	Bottom Depth	Description	Purpose
RB1	431936 E 7565566 N	42 m	100 m away from the diffuser	Exposure site located as close to the proposed diffuser as safely and logistically possible
RB2	432304 E 7567343 N	71 m	~2 km seaward of the proposed diffuser, in the center of Roberts Bay	Central monitoring site on seaward side of proposed diffuser; midway between southern shore of Roberts Bay and Melville Sound; 2 stations are already in place in the nearshore environment
REF-Marine 2	441984 E 7565159 N	42 m	~2.3 km seaward from existing Reference Site (REF-Marine 1)	To provide a reference station for the 2 new proposed AEMP stations in Roberts Bay, particularly Station RB1

Coordinates are in NAD83 UTM Z13N.

2.2 2011 SAMPLING SCHEDULE

The 2011 AEMP Marine Expansion Baseline sampling schedule paralleled that of the 2011 Doris North AEMP program. Sampling commenced in April and ended in late September. Physical oceanographic characteristics (e.g., temperature, salinity, dissolved oxygen) and water quality (e.g., nutrients and metals) were collected in Roberts Bay four times during the sampling period. These were collected not less than one month apart (whenever possible), thereby complying with MMER guidelines (Schedule 5, s.7 (1-2)) and the 2011 Doris North AEMP program. In Reference Bay, water quality samples were collected three times in 2011 (July, August, and September) and physical profiles were collected twice (July, August). Phytoplankton biomass (as chlorophyll *a*) was collected in each bay during the three open-water surveys (July, August, and September) and sediment quality and benthos samples were collected once in summer (August).

Table 2.2-1 lists the parameters measured for each marine environmental component during 2011. The under-ice, April sampling was designed to collect water when dissolved oxygen concentrations would be lowest (due to reduced photosynthesis and the absence of oxygen diffusion from the atmosphere), and nutrient and metal concentrations would be highest (due to limited biological uptake and solute extrusion during ice formation).

Table 2.2-1. Environmental Components and Sampling Time, AEMP Marine Baseline Expansion, 2011

Sampled Parameters	April	July	August	September
Water Column Structure				
Salinity	X	X	X	X
Temperature	X	X	X	X
Dissolved Oxygen	X	X	X	X
Euphotic Zone Depth		X	X	X
Water Quality				
Physical, Anions, and Nutrients	X	X	X	X
Metals (Total)	X	X	X	X
Sediment Quality				
Particle Composition, % Moisture			X	
Organic Content, Nutrients			X	
Total Metals			X	
Biology				
Phytoplankton Biomass	X	X	X	X
Benthos Abundance, Taxonomy, and Diversity			X	

2.3 SAMPLING METHODS

Table 2.3-1 provides a summary of sampled parameters, replication, sampling dates, and sampling equipment used. Full descriptions are provided in the following text.

2.3.1 Physical Oceanography and Light Penetration

In Roberts Bay, physical oceanographic profiles (e.g., salinity, temperature, and dissolved oxygen) were collected once during the under-ice season (April) and three times during the open-water season (July, August, and September) in 2011 (Table 2.3-2). Physical profiles were collected in Reference Bay in July and August.

In April, the under-ice water was accessed by drilling a 25-cm diameter hole through the ice. The ice thickness was recorded and a vertical profile of temperature, salinity, and conductivity was collected using an *in situ* conductivity, temperature, and depth probe (CTD; RBR-420). The probe was lowered through the water column at an approximate speed 0.5 m/s to within 1 m of the sea floor. The data logged during this process were immediately transferred to a computer in the field. Data from the downcast were used to derive physical profiles. Following the CTD casts, vertical profiles of dissolved oxygen concentration and percent saturation were collected using a YSI dissolved oxygen meter lowered at 1 m intervals to within 1 m of the sediments.

During the open-water surveys (July-September), CTD and dissolved oxygen profiles were collected from the side of an aluminum boat. The logged and recorded data were processed using the same methods in April.




Figure 2.1-1

Marine AEMP Expansion Sampling Locations, Doris North Project, 2011

Table 2.3-1. Summary of the Marine AEMP Expansion Baseline Sampling Program, Doris North Project, 2011

Monitoring Parameter	Sampling Frequency	Sample Replication and Depths	Sampling Dates/Timing	Sampling Device
Physical Oceanography				
Temperature, salinity, and dissolved oxygen profiles, Secchi depth	4 x	Entire water column; n = 1/site	April, July, August, September	Conductivity-Temperature-Depth (CTD) probe; dissolved oxygen meter
Water Quality				
Physical, nutrients, cyanides, total metals, radium 226	4 x	4 depths; n = 1/depth @ 1 m below the surface, 4 m above pycnocline, 4 m below pycnocline, mid-deep layer + 20% replication	April, July, August, September	2.5 L Niskin (April); 5 L GO-FLO sampling bottle (July-Sept)
Sediment Quality				
Physical, particle size, nutrients, metals, total organic carbon (TOC)	1 x	n = 3/site	August; coincident with August marine survey	Petite Ponar grab
Phytoplankton Biomass				
Chlorophyll <i>a</i>	3 x	n = 3/site @ 1 m below the surface	July, August, September; coincident with water quality surveys	2.5 L Niskin (April); 5 L GO-FLO sampling bottle (July-Sept)
Benthos				
Abundance and taxonomy	1 x	n = 5/site (3 subsamples/replicate)	August; coincident with August survey	Petite Ponar grab; 500 µm sieve

Table 2.3-2. 2011 Physical Oceanography Sampling Dates, Marine AEMP Expansion Program

Site	CTD and DO Sampling Date	Secchi Depth Sampling Date
RB1 (Roberts Bay)	23-Apr-11	n/c
	21-Jul-11	22-Jul-11
	14-Aug-11	14-Aug-11
	22-Sep-11	22-Sep-11
RB2 (Roberts Bay)	23-Apr-11	n/c
	22-Jul-11	22-Jul-11
	14-Aug-11	14-Aug-11
	22-Sep-11	22-Sep-11
REF-Marine 2 (Reference Bay)	n/c	n/c
	27-Jul-11	27-Jul-11
	19-Aug-11	19-Aug-11
	n/c	25-Sep-11

DO - dissolved oxygen; n/c - not collected

Secchi depths were measured during July, August, and September to determine the light penetration at each site during the open-water season. A 30-cm white disk was placed over the shaded side of the boat and lowered until it disappeared from sight. It was raised into view and lowered until it disappeared a second time. This was recorded as the Secchi depth (D_s). The 1% ($Z_{1\%}$) euphotic zone depth was computed by first calculating the light extinction coefficient (k) from D_s , and then

calculating the euphotic zone depth based on the appropriate light extinction coefficient. The 1% euphotic depth is the depth of the water column where 1% of the surface irradiance reaches. It represents the depth at which the integrated gross water column photosynthetic production is equivalent to the integrated gross water column respiration; thus, there is net photosynthesis above this depth. The 1% euphotic depth is often known as the compensation depth, and is calculated as follows (Parsons et al. 1984):

Light extinction coefficient:

$$k \text{ (m}^{-1}\text{)} = 1.7/D_s$$

Euphotic Depth (1%):

$$Z_{1\%} \text{ (m)} = 4.6/k$$

2.3.2 Water Quality

Water quality samples were collected at RB1 and RB2 during the under-ice season in April, and at all three AEMP Expansion sites during the open-water season in July, August, and September 2011. As much as possible, samples at a specific site were collected more than one month apart to conform to EEM recommendations (Environment Canada 2002a). Depths of water quality sampling were determined based on the water column structure (as determined by the CTD profiles initially collected at a specific site). At each site, four depths were sampled: 1 m below the surface, 4 m above the pycnocline, 4 m below the pycnocline, and at mid-depth of the deep layer. The pycnocline was defined as the depth zone where the density changed most sharply. Ultimately, the four depths were distributed to best approximate the water column structure as determined by the CTD profiles. The sampling dates and depths for all water quality sampling are presented in Table 2.3-3 and the analyzed parameters are summarized in Table 2.3-4. All sampling locations are presented in Figure 2.1-1. The sampling procedures used are described below.

Table 2.3-3. 2011 Water Quality Sampling Dates and Depths, Marine AEMP Expansion Program

Site	Sampling Date	Depth (m)
RB1 (Roberts Bay)	24-Apr-11*	1, 25, 33, 48
	22-Jul-11	1, 7, 15, 40
	14-Aug-11	1, 6, 14, 35
	22-Sep-11	1, 8, 21, 40
RB2 (Roberts Bay)	23-Apr-11*	1, 35, 44, 48
	22-Jul-11	1, 7, 15, 40
	14-Aug-11	1, 7, 15, 40
	22-Sep-11	1, 9, 21, 40
REF-Marine 2 (Reference Bay)	27-Jul-11	1, 8, 20, 38
	19-Aug-11	1, 6, 14, 40
	25-Sep-11	1, 8, 20, 40

* April surface water quality samples were collected from 1 m below the ice.

In April, the underlying water was accessed through a hole following the temperature, CTD, and dissolved oxygen profiles. An adapted 2.5 L ‘skinny’ Niskin bottle was used to collect water during winter sampling. This bottle is designed to ‘trip’ and collect discrete samples during freezing temperatures. To avoid metal contamination, the tripping mechanism used acid-cleaned silicone tubing within the interior of the bottle. A dual rope system was used to achieve bottle closure and to ensure the collection of discrete samples. During the summer and fall sampling, all water samples were collected using an acid-washed 5 L GO-FLO sampling device. The GO-FLO was securely attached to a metred cable line and was then

lowered to the appropriate sampling depth. It was allowed to equilibrate for one minute, at which time it was triggered closed using a brass messenger. The GO-FLO was then raised and brought into the boat for subsampling. Each GO-FLO cast represented one replicate.

Table 2.3-4. Marine AEMP Expansion Water Quality Parameters and Realized Detection Limits, Doris North Project, 2011

Parameter	Units	Realized Detection Limits	Parameter	Units	Realized Detection Limits
Physical Tests			Total Metals (cont'd)		
Hardness (as CaCO ₃)	mg/L	4.3	Copper (Cu)	mg/L	0.000050
pH	pH	0.10	Gallium (Ga)	mg/L	0.00050
Salinity (Electrical Conductivity)	g/L	1.0	Iron (Fe)	mg/L	0.010
Total Suspended Solids	mg/L	2.0 to 4.0	Lead (Pb)	mg/L	0.000050
Turbidity	NTU	0.10	Lithium (Li)	mg/L	0.020
Anions			Magnesium (Mg)	mg/L	1.0
Alkalinity, Total (as CaCO ₃)	mg/L	2.0	Manganese (Mn)	mg/L	0.000050
Bromide (Br)	mg/L	5.0	Mercury (Hg)	mg/L	0.000010
Chloride (Cl)	mg/L	50	Mercury (Hg) (ultra low-level)	µg/L	0.00050
Fluoride (F)	mg/L	0.03 or 0.75	Molybdenum (Mo)	mg/L	0.0020
Sulphate (SO ₄)	mg/L	50	Nickel (Ni)	mg/L	0.000050
Nutrients			Phosphorus (P)	mg/L	1.0
Ammonia (as N)	mg/L	0.0050	Potassium (K)	mg/L	20
Nitrate and Nitrite (as N)	mg/L	0.0060	Rhenium (Re)	mg/L	0.00050
Nitrate (as N)	mg/L	0.0060	Rubidium (Rb)	mg/L	0.0050
Nitrite (as N)	mg/L	0.0020	Selenium (Se)	mg/L	0.00040
Total Phosphorus	mg/L	0.0020	Silicon (Si)	mg/L	0.50
Cyanides			Silver (Ag)	mg/L	0.00010 or 0.00020
Total Cyanide	mg/L	0.0010	Sodium (Na)	mg/L	20
Free Cyanide	mg/L	0.0010 or 0.0050	Strontium (Sr)	mg/L	0.010 or 0.050
Organics			Tellurium (Te)	mg/L	0.00050
Total Organic Carbon	mg/L	0.50	Thallium (Tl)	mg/L	0.000050
Total Metals			Thorium (Th)	mg/L	0.00050
Aluminum (Al)	mg/L	0.0050	Tin (Sn)	mg/L	0.0010 or 0.0020
Antimony (Sb)	mg/L	0.00050	Titanium (Ti)	mg/L	0.0050
Arsenic (As)	mg/L	0.0004	Tungsten (W)	mg/L	0.0010
Barium (Ba)	mg/L	0.0010	Uranium (U)	mg/L	0.000050
Beryllium (Be)	mg/L	0.00050	Vanadium (V)	mg/L	0.00050
Bismuth (Bi)	mg/L	0.00050	Yttrium (Y)	mg/L	0.00050
Boron (B)	mg/L	0.10	Zinc (Zn)	mg/L	0.00080
Cadmium (Cd)	mg/L	0.000020	Zirconium (Zr)	mg/L	0.00050
Calcium (Ca)	mg/L	0.50	Speciated Metals		
Cesium (Cs)	mg/L	0.00050	Methylmercury (Total)	µg/L	0.000050
Chromium (Cr)	mg/L	0.00010 to 0.00050	Radiochemistry		
Cobalt (Co)	mg/L	0.000050	Radium-226	Bq/L	0.005

Subsamples for the various water quality parameters (e.g., physical parameters/major ions, nutrients, and total metals) were drawn from the GO-FLO/Niskin bottles. All sample bottles were rinsed three times prior to filling with care being taken not to bring the bottle or cap into contact with the plastic spigot or other possible sources of contamination. The appropriate preservatives provided by the analytical laboratory (ALS Laboratory Group, Burnaby, British Columbia) were added in the field after collection.

All samples were kept cold and in the dark while in the field and were refrigerated at camp prior to transport. Samples were transported in coolers with freezer packs to ALS, Burnaby for all analyses. The parameters analyzed and applicable detection limits are summarized in Table 2.3-4.

All water quality parameters were screened against the Canadian Council of the Ministers of the Environment (CCME) water quality guidelines for the protection of aquatic life (CCME 2011a).

2.3.3 Sediment Quality

Sediment quality samples were collected at RB1 (Roberts Bay) and REF-Marine 2 (Reference Bay) during the open-water season in August 2011. This sampling coincided with benthic invertebrate sampling. The sampling dates and depths for each of the two sites are presented in Table 2.3-5 and the analyzed parameters are summarized in Table 2.3-6. All sampling locations are presented in Figure 2.1-1. The sampling procedures are described below.

Table 2.3-5. 2011 Sediment Quality and Benthic Invertebrate Sampling Dates and Depths, Marine AEMP Expansion Program

Site	Sampling Date	Depth (m)
RB1 (Roberts Bay)	23-Aug-10	42
REF-Marine 2 (Reference Bay)	24-Aug-10	42

Table 2.3-6. Marine AEMP Expansion Sediment Quality Parameters and Realized Detection Limits, Doris North Project, August 2011

Parameter	Units	Realized Detection Limits	Parameter	Units	Realized Detection Limits
Moisture	%	0.25	Metals (cont'd)		
pH	pH	0.10	Cobalt (Co)	mg/kg	0.10
Particle Size			Copper (Cu)	mg/kg	0.50
% Gravel (>2 mm)	%	0.10	Iron (Fe)	mg/kg	50
% Sand (2.0 mm - 0.063 mm)	%	0.10	Lead (Pb)	mg/kg	0.50
% Silt (0.063 mm - 4 µm)	%	0.10	Lithium (Li)	mg/kg	1.0
% Clay (<4 µm)	%	0.10	Magnesium (Mg)	mg/kg	20
Nutrients			Manganese (Mn)	mg/kg	1.0
Total Nitrogen	%	0.020	Mercury (Hg)	mg/kg	0.0050 or 0.015
Organic / Inorganic Carbon			Molybdenum (Mo)	mg/kg	0.50
Total Organic Carbon	%	0.10	Nickel (Ni)	mg/kg	0.50
Plant Available Nutrients			Phosphorus (P)	mg/kg	50
Available Ammonium-N	mg/kg	1.6 or 2.4	Potassium (K)	mg/kg	100
Available Nitrate-N	mg/kg	4.0 or 6.0	Selenium (Se)	mg/kg	0.20
Available Nitrite-N	mg/kg	0.80 or 1.2	Silver (Ag)	mg/kg	0.10

(continued)

Table 2.3-6. Marine AEMP Expansion Sediment Quality Parameters and Realized Detection Limits, Doris North Project, August 2011 (completed)

Parameter	Units	Realized Detection Limits	Parameter	Units	Realized Detection Limits
Available Phosphate-P	mg/kg	4.0 or 6.0	Sodium (Na)	mg/kg	100
Metals			Strontium (Sr)	mg/kg	0.50
Aluminum (Al)	mg/kg	50	Sulphur (S)	mg/kg	500
Antimony (Sb)	mg/kg	0.10	Thallium (Tl)	mg/kg	0.050
Arsenic (As)	mg/kg	0.050	Tin (Sn)	mg/kg	2.0
Barium (Ba)	mg/kg	0.50	Titanium (Ti)	mg/kg	1.0
Beryllium (Be)	mg/kg	0.20	Uranium (U)	mg/kg	0.050
Bismuth (Bi)	mg/kg	0.20	Vanadium (V)	mg/kg	0.20
Cadmium (Cd)	mg/kg	0.050	Zinc (Zn)	mg/kg	1.0
Calcium (Ca)	mg/kg	50	Speciated Metals		
Chromium (Cr)	mg/kg	0.50	Methylmercury	mg/kg	0.000050

Marine sediment quality samples were collected in triplicate using a Petite Ponar grab sampler, with replicates spaced 20 to 50 m apart. Each sediment sample was carefully transferred onto a white, plastic tray, and the top 2 cm of sediment was scraped into a bowl, homogenized, and then placed into two Whirl-Pak bags: one for grain size, and one for sediment chemistry. All samples were refrigerated (in darkness) until they were shipped to ALS in Yellowknife, NT, for analysis on the first available flight out of Doris camp.

All sediment quality parameters were screened against the CCME quality guidelines for the protection of aquatic life (CCME 2011b).

2.3.4 Phytoplankton

Phytoplankton biomass (as chlorophyll *a*) samples were collected at RB1 and RB2 in Roberts Bay during April, July, August, and late September. Samples were collected at REF-Marine 2 in Reference Bay in July, August, and late September. Triplicate samples were collected at 1 m depth using a 5 L GO-FLO water sampler. All samples were transferred into 1 L plastic bottles and stored in a cool, dark cooler until being returned to camp later in the day. Once at camp, the samples were filtered onto 0.45 µm filters, which were wrapped into aluminum foil, and stored frozen. Chlorophyll samples were hand carried to Vancouver, BC, to ensure they remained frozen and sent to ALS (Burnaby, BC) for analysis. The sampling dates are presented in Table 2.3-7 and all sampling locations are presented in Figure 2.1-1.

2.3.5 Benthos

Benthic invertebrate samples were collected during summer at the shallower AEMP site in Roberts Bay (RB1) and at the reference site in Reference Bay (REF-Marine 2). This coincided with the sediment quality sampling. All field sampling devices and methods for the AEMP Expansion benthos sampling were designed to comply with EEM guidance documents (Environment Canada 2002a). Sampling dates and depths for each of the two sites are presented in Table 2.3-5 and all sampling locations are indicated in Figure 2.1-1.

Samples were collected with a Petite Ponar dredge. To comply with EEM requirements, three separate subsamples were collected and pooled for each replicate sample, and five replicates were collected at each site. Each replicate was collected approximately 20 to 50 m apart. The Petite Ponar was carefully set

open, lowered gradually onto the sediment using a metred cable line, and triggered closed. Once at the surface, each sediment sample was transferred into a 500 µm sieve bucket and rinsed with site-specific water until free of sediments. The material retained within the sieve was then placed into a labelled plastic jar and filled with 10% buffered formalin. All benthos samples were sent to Columbia Science (Courtney, BC) for enumeration and identification, and community descriptors were calculated as described below. Benthos density was calculated by dividing the total benthos counts by the surface area of the of the Petite Ponar sampler (0.023 m²).

Table 2.3-7. 2011 Phytoplankton Biomass Sampling Dates and Depths, Marine AEMP Expansion Program

Site	Sampling Date
RB1 (Roberts Bay)	23-Apr-11
	23-Jul-11
	14-Aug-11
	22-Sep-11
RB2 (Roberts Bay)	23-Apr-11
	23-Jul-11
	14-Aug-11
	22-Sep-11
REF-Marine 2 (Reference Bay)	21-Jul-11
	19-Aug-11
	25-Sep-11

Several community descriptors were calculated from the taxonomic results, including benthos density, family richness, Simpson's Diversity and Evenness indices, and the Bray-Curtis Index. Cyclopoid and calanoid copepods were not included in the community metrics as they are generally planktonic. Nematodes and harpacticoid copepods were excluded as they are considered meiofauna (invertebrates between 63 µm and 500 µm) and were not adequately sampled using a 500 µm sieve bucket. Those organisms that could not be identified to at least the family level, and made up less than 2% of the benthos counts in each replicate, were also excluded from the community analysis. Family richness was calculated as the total number of benthic invertebrate families present in each replicate sample, and the Simpson diversity index was calculated as:

$$\text{Simpson Diversity Index (D)} = 1 - \sum_{i=1}^F p_i^2$$

where F is the number of families present (i.e., family richness), and p_i is the relative abundance of each family calculated as n_i/N , where n_i is the number of individuals in family i , and N is the total number of all individuals. Simpson's Evenness Index was calculated as:

$$\text{Simpson Evenness Index (E)} = 1 / \sum_{i=1}^F p_i^2 / F$$

A complete dissimilarity matrix was also generated that included pair-wise comparisons of all samples using the Bray-Curtis Index (Environment Canada 2002a). This index takes into account the number of families and the total number of individuals within each replicate sample and compares it to a single median reference community composition. This reference composition was generated from the median abundance of each represented family from all of the reference site replicates. Since the median reference composition was generated from the reference site replicates, the comparison of an actual

reference site replicate community to the median reference community composition will produce a dissimilarity value (although generally a much lower value than exposure sites). Because the Bray-Curtis Index measures the percent difference between sites, the greater the dissimilarity value between a site and the median reference community, the more dissimilar those benthos communities are. This index is calculated as:

$$\text{Bray-Curtis Index (BC)} = \sum_{i=1}^n |y_{i1} - y_{i2}| / \sum_{i=1}^n (y_{i1} + y_{i2})$$

where BC is the Bray-Curtis distance between sites 1 and 2, n is the total number of families present at the two sites, y_{i1} is the count for family i at site 1, and y_{i2} is the count for family i at site 2.

2.3.6 Quality Assurance and Quality Control (QA/QC)

A quality assurance and quality control (QA/QC) program was incorporated into the design of this study to track samples, account for within site variability, and identify potential sources of contamination from sampling equipment, sample transportation, or sample processing. This included the use of chain of custody forms, replicate samples, and sample blanks.

2.3.6.1 Water Quality QA/QC

All water quality samples were recorded on chain of custody forms before being sent to ALS laboratories. Replicate samples accounted for approximately 20% of water samples collected during each sampling period. These replicate samples were collected from multiple depths.

Equipment, field and travel blanks (13% of total samples) were processed and submitted with the water samples as part of the QA/QC program. These blanks identified potential sources of contamination to the field samples.

Equipment blanks were designed to identify possible contamination of the sampling equipment, including the sampling bottle (GO-FLO or Niskin). These were collected in the field by first thoroughly rinsing an acid-washed GO-FLO with double de-ionized water (provided by ALS), then filling the GO-FLO bottle with double de-ionized water. The water remained in the GO-FLO for one minute (as would occur with a real sample) until sub-samples were drawn from the sampling bottle. Equipment blanks were preserved and handled in the same manner as water quality samples.

Field blanks were designed to identify possible contamination from the exposure of the sample to ambient conditions at the sampling site. These were processed in the field by opening the bottles provided by ALS (containing double de-ionized water) and exposing the sample to ambient air for one minute. The bottles were preserved and handled in the same manner as water quality samples.

Travel blanks were designed to identify potential contamination from the transportation of samples from the sampling location to ALS laboratories. These bottles contained double de-ionized water and were provided by ALS. They were never opened in the field, and were returned to ALS with the other water quality samples. The blanks were analyzed for the same parameters as the regular water samples.

2.3.6.2 Sediment Quality QA/QC

The sediment quality QA/QC program included the use of chain of custody forms for all samples, and the collection of triplicate sediment samples at all sites to account for within-site variability.

2.3.6.3 Aquatic Biology QA/QC

Chain of custody forms were used for all aquatic biology samples. All samples had replication; triplicate samples were taken for chlorophyll *a* analysis (phytoplankton biomass), and five replicates of three composite samples were collected for benthos. Additional QA/QC measures were used by the benthos taxonomist to ensure consistent and accurate sorting of benthos samples. These procedures are outlined below.

Benthos Sorting QA/QC

As part of the QC program, the re-sorting of benthic sample residues was conducted on a randomly selected 10% of the benthos samples to determine the level of sorting efficiency. The criterion for an acceptable sorting was that more than 90% of the total number of organisms found in both the initial and QA/QC sort were recovered during the initial sort, as required by Environment Canada for invertebrate community surveys (Environment Canada, 2002b). This was calculated by the following equation:

$$\% \text{ Sorting Efficiency} = [1 - (\# \text{ in QA/QC re-sort} / (\# \text{ sorted originally} + \# \text{ QA/QC resort}))] \times 100$$

Any sample not meeting the 90% removal criterion was re-sorted a third time. The 90% minimum efficiency was attained for all benthos samples in this study.

DORIS NORTH GOLD MINE PROJECT
2011 Aquatic Effects Monitoring Program (AEMP)
Marine Expansion Baseline Report

3. Results

3. Results

3.1 PHYSICAL OCEANOGRAPHY AND LIGHT PENETRATION

Figures 3.1-1 to 3.1-3 provide graphical results of temperature, salinity, and dissolved oxygen. Table 3.1-1 provides a summary of measured Secchi depths, euphotic zone depths, and pycnocline depths. All raw data are provided in appendices, and a summary of the results is provided below.

Table 3.1-1. Marine AEMP Expansion Secchi Depths, Euphotic Zone Depths, and Estimated Pycnocline Depths, Doris North Project, 2011


Site	Depth (m)	Secchi Depth (m)			Euphotic Zone Depth (m)			Pycnocline Depth (m)*		
		July	August	September	July	August	September	July	August	September
<i>Roberts Bay</i>										
RB1	42	7	11.3	11	18.9	30.6	29.8	6.0	10.2	17.4
RB2	71	7.9	14.3	10.5	21.4	38.7	28.4	6.6	7.2	22.2
<i>Reference Bay</i>										
REF-Marine 2	42	7.1	6.6	6	19.2	17.9	16.2	5.2	3.0	26.9


* Pycnocline depths are estimates where the deepest portion of the surface density gradient was selected.


In April, the Roberts Bay water column was weakly stratified below the snow and ice, with a deep mixed layer extending to 30 m. The surface salinity was near 26.8, and increased little to 27.1 below 30 m. April water temperatures were approximately -1.4°C to -1.3°C in the upper 30 m and -0.60°C to -0.25°C in bottom waters. Dissolved oxygen concentrations reached 11.9 mg/L just below the sea ice and declined steadily to 9.2 mg/L in deeper waters.

During the summer, vertical stratification intensified in Roberts and Reference bays due to melting sea ice and increased riverine inputs. The influx of freshwater resulted in shallower pycnoclines (range: 3-10 m) with warmer, fresher water (lower density) sitting above colder, saltier (higher density) water at depth. The surface layer salinities were between 23 and 24 in July and August, with deep water salinities near 27.1. Temperatures were warmest in the surface water during August, reaching 10°C. Dissolved oxygen concentrations were lowest during this month, with a maximum surface concentration of 9.53 mg/L in Robert Bay, and a minimum deep water concentration of 6.63 mg/L in Reference Bay (41 m). This was below the recommended CCME marine water quality guideline for the protection of aquatic life (8.0 mg/L). The dissolved oxygen minimum in Roberts Bay was 8.24 mg/L at 60 m and was within the natural range for healthy coastal waters.

The strong summer stratification lasted approximately two months before the higher winds and lower temperatures in September began to mix the stable surface layer. Very strong winds were present during the September sampling and Roberts and Reference bays were laterally well mixed, with vertical mixed layer depths reaching 20 m in Roberts Bay and extending to 25 m in Reference Bay. Surface salinities increased during this period, ranging from 25.6 to 25.8. The surface temperatures ranged from 3.2°C to 3.8°C. Deep water temperatures and salinities were very similar to those in the winter and summer, approaching -0.6°C and 27.4. Dissolved oxygen concentrations reached 11.7 mg/L in the surface waters of Roberts Bay, and were 10.2 mg/L at depth.







Light penetration was high during the summer and fall in both Roberts and Reference bays. The euphotic depth (1% light level) was generally deeper than 16 m and usually extended far below the pycnocline. This suggests that net photosynthesis was possible throughout the upper water column in both bays.

3.2 WATER QUALITY

All water quality parameters with CCME guidelines are presented graphically in Figures 3.2-1 to 3.2-10 and all raw water quality data and QA/QC results are in Appendices 3.2-1 and 3.2-2. All water quality parameters with CCME guidelines are presented graphically in Figures 3.2-1 to 3.2-10 and all raw water quality data are in Appendix 3.2-1.

Overall, the water quality parameters were similar between Roberts Bay and Reference Bay. pH averaged 7.8 across all sites and depths in winter and 7.9 in summer. Water clarity was high in both bays as total suspended solids (TSS) and turbidity levels were consistently low (mean: 0.46 NTU). Some nutrients showed vertical concentration gradients, which were most dramatic during July and August. Nitrate and total phosphorus concentrations were generally lower at the surface than in deep waters. During summer, nitrate was usually undetectable (>0.006 mg N/L) in the surface layer, while total phosphorus was detectable throughout the water column, averaging 0.038 mg/L over all sites and depths. Thus, phytoplankton growth was likely controlled by nitrogen availability when sufficient light was present, which is typical of northern, estuarine systems. Vertical gradients of total organic carbon (TOC) exhibited the opposite trend as nutrients as the summer surface concentrations were usually slightly higher than at depth. Vertical gradients characterized by lower nutrients and higher TOC at the surface than at depth are indicative of phytoplankton growth and biological nutrient uptake.


Roberts Bay and Reference Bay can be considered low-metal environments, as all water quality parameters were below their respective CCME guideline for the protection of marine aquatic life. Total aluminum and iron were below detection at all depths in April and often above detection during the open-water season indicating that aerial and riverine transport were responsible for delivering these metals into Roberts and Reference bays. Cadmium, molybdenum, and nickel tended have lower concentrations in the surface waters than at depth, suggesting biological uptake or dilution from freshwater inputs. Trace amounts of radium-226 were found at all sites during most seasons, except in southern Roberts Bay in August and September. The highest concentration occurred in Reference Bay in July (0.02 mg/L at 20 m). Total lead and zinc concentrations were usually below analytical detection limits.

Total cyanide was commonly observed at low concentrations during the surveys. The highest concentrations were found in July in the surface waters of Roberts (0.0021 mg/L) and Reference (0.0019 mg/L) bays, and detectable levels were found throughout the Roberts Bay water column in April. Free cyanide made up a very small proportion of the total cyanide pool as there was little detectable free cyanide in either bay.

3.3 SEDIMENT QUALITY

Graphical results of sediment quality data are provided in Figures 3.3-1 through 3.3-3. All raw data are provided in appendices, and a summary of the results is provided below.

All sediment quality parameters were screened against the CCME sediment quality guidelines for the protection of aquatic life (CCME 2011b). Summaries of CCME exceedances can be found in Tables 3.3-1 and 3.3-2.



Notes: Error bars represent the standard error of the mean.


Black dotted lines represent the analytical detection limits.

Red dashed lines represent the CCME marine and estuarine guideline pH range (7.0–8.7).

pH is a required parameter for water quality monitoring as per Schedule 5, s. 7(1)(c) of the MMR.

Total alkalinity is a required parameter for effluent characterization and water quality monitoring as per Schedule 5, s. 4(1) and s. 7(1)(c) of the MMR.

Figure 3.2-1



Notes: Error bars represent the standard error of the mean.


Black dotted lines represent the analytical detection limits; values below the detection limit are plotted at half the applicable detection limit.

The CCME marine guideline for total suspended solids is dependent upon background levels.

Hardness is a required parameter for effluent characterization and water quality monitoring as per Schedule 5, s. 4(1) and s. 7(1)(c) of the MMER.

Total suspended solids are regulated as deleterious substances in effluents as per Schedule 4 of the MMER.

Figure 3.2-2




Notes: Error bars represent the standard error of the mean.

Black dotted lines represent the analytical detection limits; values below the detection limit are plotted at half the applicable detection limit.

Red dashed line represents the interim CCME marine guideline for nitrate as N (3.612 mg/L).

Total ammonia and nitrate are required parameters for effluent characterization as per Schedule 5, s. 4(1) of the MMER.

Figure 3.2-3




Notes: Error bars represent the standard error of the mean.

Black dotted lines represent the analytical detection limits; values below the detection limit are plotted at half the applicable detection limit.

Total cyanide is regulated as a deleterious substance in effluents as per Schedule 4 of the MMER.

Figure 3.2-4




Notes: Error bars represent the standard error of the mean.

Black dotted lines represent the analytical detection limits; values below the detection limit are plotted at half the applicable detection limit.

Radium-226 is regulated as a deleterious substance in effluents as per Schedule 4 of the MMER.

Total aluminum is a required parameter for effluent characterization as per Schedule 5, s. 4(1) of the MMER.

Figure 3.2-5



Notes: Error bars represent the standard error of the mean.


Black dotted lines represent the analytical detection limits; values below the detection limit are plotted at half the applicable detection limit.

Red dashed lines represent the interim CCME marine guideline for arsenic (0.0125 mg/L) and the CCME marine guideline for cadmium (0.00012 mg/L).

Total arsenic is regulated as a deleterious substance in effluents as per Schedule 4 of the MMER.

Total cadmium is a required parameter for effluent characterization as per Schedule 5, s. 4(1) of the MMER.

Figure 3.2-6




Notes: Error bars represent the standard error of the mean.

Black dotted lines represent the analytical detection limits; values below the detection limit are plotted at half the applicable detection limit.

Total copper is regulated as a deleterious substance in effluents as per Schedule 4 of the MMER.

Total iron is a required parameter for effluent characterization as per Schedule 5, s. 4(1) of the MMER.

Figure 3.2-7




Notes: Error bars represent the standard error of the mean.

Black dotted lines represent the analytical detection limits; values below the detection limit are plotted at half the applicable detection limit.

Total lead is regulated as a deleterious substance in effluents as per Schedule 4 of the MMER.

Total molybdenum is a required parameter for effluent characterization as per Schedule 5, s. 4(1) of the MMER.

Figure 3.2-8




Notes: Error bars represent the standard error of the mean.

Black dotted lines represent the analytical detection limits; values below the detection limit are plotted at half the applicable detection limit.

Red dashed line represents the interim CCME marine guideline for inorganic mercury (0.000016 mg/L).

Total mercury is a required parameter for effluent characterization as per Schedule 5, s. 4(1) of the MMER.

Figure 3.2-9




Notes: Error bars represent the standard error of the mean.

Black dotted lines represent the analytical detection limits; values below the detection limit are plotted at half the applicable detection limit.


Total nickel and total zinc are regulated as deleterious substances in effluents as per Schedule 4 of the MMER.

Figure 3.2-10



Notes: Error bars represent the standard error of the mean.
Stacked bars represent the mean of replicate samples.
Black dotted line represents the analytical detection limit;
values below the detection limit are plotted at half the detection limit.

Figure 3.3-1



Notes: Error bars represent the standard error of the mean.


Black dotted lines represent analytical detection limits; values below the detection limit are plotted at half the detection limit.

Red dashed lines represent the CCME marine and estuarine interim sediment quality guidelines (ISQGs) for arsenic (7.24 mg/kg), cadmium (0.7 mg/kg), chromium (52.3 mg/kg), and copper (18.7 mg/kg), and the probable effects level (PEL) for arsenic (41.6 mg/kg).

The PELs for cadmium (4.2 mg/kg), chromium (160 mg/kg), and copper (108 mg/kg) are not shown.

**Total Arsenic, Total Cadmium, Total Chromium,
and Total Copper Concentrations in Sediments
at Marine AEMP Expansion Sites, 2011**

Figure 3.3-2



Notes: Error bars represent the standard error of the mean.

Black dotted lines represent analytical detection limits; values below the detection limit are plotted at half the detection limit.

Red dashed lines represent the CCME marine and estuarine interim sediment quality guidelines (ISQGs) for total mercury (0.13 mg/kg), lead (30.2 mg/kg) and zinc (124 mg/kg); probable effects levels (PELs) for total mercury (0.7 mg/kg), lead (112 mg/kg), and zinc (271 mg/kg) are not shown.

Table 3.3-1. Sediment Quality at Marine AEMP Expansion Sites, Percent of Samples in which Concentrations are Higher than CCME Guidelines, 2011

Marine Site	Total Number of Samples Collected	CCME Guideline value ^a (mg/kg):	Percent of samples higher than ISQG ^b guidelines					
			Arsenic (As)	Cadmium (Cd)	Chromium (Cr)	Copper (Cu)	Lead (Pb)	Zinc (Zn)
			7.24	0.7	52.3	18.7	30.2	124
RB1	3		100	0	100	100	0	0
REF-Marine 2	3		100	0	0	66.7	0	0
Marine Site	Total Number of Samples Collected	CCME Guideline value ^a (mg/kg):	Percent of samples higher than PEL ^c guidelines					
			Arsenic (As)	Cadmium (Cd)	Chromium (Cr)	Copper (Cu)	Lead (Pb)	Zinc (Zn)
			41.6	4.2	160	108	112	271
RB1	3		66.7	0	0	0	0	0
REF-Marine 2	3		100	0	0	0	0	0

Values represent the percentages of 2011 samples that are higher than CCME guidelines.

a) Canadian Sediment Quality Guidelines for the Protection of Freshwater Aquatic Life, CCME, updated January 2011.

b) ISQG = Interim Sediment Quality Guideline.

c) PEL = Probable Effects Level.

Table 3.3-2. Sediment Quality at Marine AEMP Expansion Sites, Factor by which Average Concentrations are Higher than CCME Guidelines, 2011

Marine Site	Total Number of Samples Collected	CCME Guideline value ^a (mg/kg):	Factor by which average concentrations are higher than ISQG ^b guidelines					
			Arsenic (As)	Cadmium (Cd)	Chromium (Cr)	Copper (Cu)	Lead (Pb)	Zinc (Zn)
			7.24	0.7	52.3	18.7	30.2	124
RB1	3		5.09	-	1.24	1.32	-	-
REF-Marine 2	3		7.09	-	-	1.02	-	-
Marine Site	Total Number of Samples Collected	CCME Guideline value ^a (mg/kg):	Factor by which average concentrations are higher than PEL ^c guidelines					
			Arsenic (As)	Cadmium (Cd)	Chromium (Cr)	Copper (Cu)	Lead (Pb)	Zinc (Zn)
			41.6	4.2	160	108	112	271
RB1	3		-	-	-	-	-	-
REF-Marine 2	3		1.23	-	-	-	-	-

The average 2011 concentration of a particular parameter may be below the CCME guideline concentration even if a percentage of samples is higher than this guideline concentration.

a) Canadian Sediment Quality Guidelines for the Protection of Freshwater Aquatic Life, CCME, updated January 2011.

b) ISQG = Interim Sediment Quality Guideline.

c) PEL = Probable Effects Level.

The deep water sediments (42 m) were composed mainly of silt (mean: 54.2%) and clay (mean: 41.3%) in both bays. Roberts Bay sediments had slightly higher mean concentrations of TOC, total chromium, total copper, total lead, and total zinc than in Reference Bay. Conversely, Reference Bay sediments had higher concentrations of total arsenic. The mean arsenic, chromium, and copper levels found in Roberts Bay sediments were higher than CCME interim marine sediment quality guidelines (ISQGs) for these metals. Mean copper levels in Reference Bay were also higher than the ISQG, and the total arsenic concentrations in this bay were particularly high, exceeding both the ISQG and the probable effects level (PEL) for the protection of marine aquatic life.

3.4 PHYTOPLANKTON BIOMASS


Figure 3.4-1 presents a graphical representation of the phytoplankton biomass (as chlorophyll *a*). All raw data are provided in appendices, and a summary of the results is provided below.

Mean phytoplankton biomass (as chlorophyll *a*) in the Project area was very low at all sites (0.14 to 0.62 µg chl *a*/L), reaching a maximum of 1.5 µg chl *a*/L in Reference Bay in August. The low concentrations reflected the low under-ice light levels in April, and the low surface layer nutrients and stable water column in July and August. In Roberts Bay, the mean phytoplankton biomass was highest in September (0.46 µg chl *a*/L), likely due to the wind-driven transport of nutrients into the surface layer during this time.


3.5 BENTHIC INVERTEBRATES


Figures 3.5-1 through 3.5-3 provide graphical results of marine benthic invertebrate data. All raw data are provided in appendices, and a summary of the results is provided below.

Mean benthos density was slightly higher in the deep Reference Bay sediments (1,520 organisms/m²) than in Roberts Bay (1,060 organisms/m²). The benthos communities were diverse in both bays (Simpson's Diversity Index: >0.80), with fewer taxonomic groups (mean richness: 15.6) being more evenly distributed (mean Simpson's Evenness: 0.49) in Roberts Bay than in Reference Bay (mean richness: 19.2; mean Simpson's Evenness: 0.30). The benthos community in Roberts Bay was largely dominated by families of polychaete worms (Maldanidae, 24.1%; Cirratulidae, 9.2%; Nephtyidae, 3.9%) and small crustaceans (Diastylidae, 15.7%; Lysianassidae, 8.6%; Ostracoda, 7.1%). The Reference Bay benthos community was more heavily dominated by polychaetes (Cirratulidae, 30.7%; Maldanidae, 19.4%; Nephtyidae, 7.8%; Capitellidae, 5.8%).




Notes: Error bars represent the standard error of the mean.
Black dotted lines represent analytical detection limits.






Notes: Error bars represent the standard error of the mean.
Stacked bars represent the mean of replicate samples.

Figure 3.5-1



Notes: Error bars represent the standard error of the mean.



Notes: Error bars represent the standard error of the mean.

DORIS NORTH GOLD MINE PROJECT
2011 Aquatic Effects Monitoring Program (AEMP)
Marine Expansion Baseline Report

References

References

- Metal Mining Effluent Regulations, SOR/2002-222. C.
- CCME. 2011a. *Canadian Water Quality Guidelines for the Protection of Aquatic Life: Summary Table. Updated January 2011. In: Canadian Environmental Quality Guidelines, 1999.* Winnipeg, MB: Canadian Council of Ministers of the Environment.
- CCME. 2011b. *Canadian Sediment Quality Guidelines for the Protection of Aquatic Life: Summary Table. Updated January 2011. In: Canadian Environmental Quality Guidelines, 1999.* Winnipeg, MB: Canadian Council of Ministers of the Environment.
- Environment Canada. 2002a. *Metal Mining Guidance Document for Aquatic Environmental Effects Monitoring.* Ottawa, ON: National Environmental Effects Monitoring Office, Environment Canada.
- Environment Canada. 2002b. *Additional Technical Guidance: Sample Sorting and Subsampling Protocols for Environmental Effects Monitoring Benthic Invertebrate Community Surveys.* <http://www.ec.gc.ca/ese-eem/default.asp?lang=En&n=B9DBF4CC-1> (accessed November 2011).
- HBML. 2011. Doris North Mine Modifications and Related Amendments to Project Certificate No. 3 and Type A Water Licence NO. 2AM-DOH0713. Prepared by Hope Bay Mining Limited for the Nunavut Water Board and the Nunavut Impact Review Board.
- Parsons, T.R., Y. Maita, and C.M. Lalli. 1984. *A Manual of Chemical and Biological Methods for Seawater Analysis.* Oxford: Pergamon Press.
- Rescan. 2010. *Doris North Gold Mine Project: Aquatic Effects Monitoring Plan.* Prepared for Hope Bay Mining Limited by Rescan Environmental Services Ltd.
- Rescan. 2011. *Doris North Gold Mine Project: 2010 Aquatic Effects Monitoring Program Report.* Prepared for Hope Bay Mining Ltd. by Rescan Environmental Services Ltd.

Appendix 3.1-1

**Temperature and Salinity Data for the Marine AEMP
Expansion Sites, Doris North Project, 2011**

Appendix 3.1-1. Temperature and Salinity Data for the AEMP Expansion Sites, Doris North Project, 2011

RB1											
23-Apr-11			21-Jul-11			14-Aug-11			22-Sep-11		
Depth (m)	Temperature (°C)	Salinity	Depth (m)	Temperature (°C)	Salinity	Depth (m)	Temperature (°C)	Salinity	Depth (m)	Temperature (°C)	Salinity
1.50	-1.40	26.84	1.05	7.48	24.79	0.97	9.76	23.61	0.68	4.06	25.41
1.88	-1.40	26.84	1.76	3.59	25.86	1.07	9.73	23.63	1.49	3.77	25.63
2.33	-1.39	26.84	2.12	3.81	25.89	1.79	9.72	23.64	1.98	3.73	25.71
2.92	-1.39	26.84	2.72	3.36	26.05	2.18	9.72	23.67	2.42	3.72	25.70
3.88	-1.40	26.85	3.13	3.37	26.13	2.54	9.75	23.68	2.74	3.67	25.75
4.81	-1.40	26.84	3.60	2.87	26.29	3.51	9.64	23.75	3.21	3.66	25.75
5.65	-1.40	26.84	4.24	2.40	26.42	4.14	9.41	23.84	3.84	3.58	25.81
5.92	-1.40	26.84	5.01	1.86	26.47	4.65	8.89	24.19	4.20	3.40	25.91
6.80	-1.39	26.85	5.53	1.16	26.54	5.38	7.75	24.58	4.74	3.29	25.96
7.77	-1.40	26.85	6.04	0.89	26.66	6.01	7.07	24.94	5.24	3.27	25.96
8.80	-1.40	26.85	6.68	0.94	26.78	6.57	6.30	25.24	5.71	3.26	25.97
9.50	-1.40	26.85	7.10	0.98	26.68	7.29	5.56	25.49	6.16	3.24	25.98
10.14	-1.40	26.84	7.52	0.79	26.73	7.88	4.51	25.58	6.77	3.21	25.99
10.81	-1.39	26.85	8.15	0.76	26.76	8.56	3.71	25.68	7.22	3.12	26.00
11.70	-1.40	26.84	8.85	0.71	26.75	9.46	3.13	25.88	7.72	3.01	26.11
11.92	-1.40	26.84	9.32	0.71	26.78	10.22	2.57	26.13	7.74	3.02	26.10
13.12	-1.39	26.84	9.78	0.71	26.77	10.94	2.25	26.21	7.70	3.02	26.10
14.25	-1.40	26.84	10.20	0.74	26.78	11.70	1.96	26.37	8.06	3.01	26.10
15.78	-1.40	26.85	11.17	0.42	26.69	12.53	1.74	26.35	8.53	3.01	26.11
16.40	-1.39	26.84	13.23	0.06	26.87	13.19	1.51	26.43	9.20	3.00	26.13
17.63	-1.39	26.85	14.06	0.01	26.84	14.01	1.29	26.51	9.76	2.98	26.14
18.51	-1.35	26.83	15.07	-0.09	26.87	14.81	1.13	26.49	10.19	2.96	26.14
19.41	-1.36	26.87	16.24	-0.15	26.87	15.54	0.93	26.60	11.12	2.90	26.17
20.09	-1.39	26.88	17.09	-0.18	26.87	16.46	0.82	26.64	12.17	2.84	26.21
21.14	-1.41	26.86	18.17	-0.25	26.87	17.25	0.74	26.65	13.17	2.84	26.22
22.29	-1.40	26.85	19.59	-0.42	26.91	18.05	0.64	26.70	14.20	2.82	26.24
23.43	-1.39	26.85	20.12	-0.43	26.92	18.85	0.57	26.72	15.23	2.82	26.24
24.51	-1.39	26.88	21.45	-0.45	26.93	19.64	0.49	26.77	16.08	2.81	26.24
25.72	-1.41	26.86	22.43	-0.46	26.94	20.54	0.44	26.76	17.42	2.62	26.28
27.22	-1.41	26.86	23.29	-0.44	26.97	21.34	0.37	26.69	18.16	2.45	26.16
28.31	-1.40	26.86	24.22	-0.41	26.95	22.20	0.24	26.74	19.04	2.07	26.26
29.26	-1.33	26.80	25.36	-0.42	26.98	22.94	0.14	26.78	20.83	1.36	26.54
30.29	-1.20	26.71	26.59	-0.41	27.01	23.76	0.07	26.80	21.62	1.12	26.64
31.56	-0.91	26.65	27.47	-0.40	27.01	24.65	-0.01	26.87	22.59	0.84	26.76
32.78	-0.42	26.56	28.11	-0.40	27.01	25.51	-0.03	26.80	23.49	0.62	26.79
33.91	-0.35	27.02	29.45	-0.40	27.01	26.39	-0.10	26.89	24.38	0.46	26.84
34.48	-0.31	26.97	30.49	-0.41	27.03	27.06	-0.13	26.87	25.51	0.35	26.87
35.19	-0.30	26.98	31.39	-0.41	27.04	28.91	-0.21	26.91	26.36	0.18	26.90
36.66	-0.31	26.99	32.08	-0.41	27.04	29.85	-0.25	26.93	27.55	-0.01	27.02
37.17	-0.31	26.99	33.27	-0.43	27.05	30.52	-0.28	26.95	28.50	-0.11	27.02
38.03	-0.31	27.01	34.36	-0.44	27.05	31.13	-0.29	26.96	29.06	-0.17	27.06
39.15	-0.34	27.03	35.22	-0.45	27.06	33.01	-0.34	26.97	30.14	-0.22	27.07
40.73	-0.31	27.00	36.30	-0.52	27.06	34.86	-0.41	27.00	31.05	-0.27	27.08
41.86	-0.33	27.01	37.76	-0.60	27.07	35.52	-0.43	27.02	32.62	-0.34	27.12
42.83	-0.31	27.01	38.76	-0.63	27.09	36.34	-0.46	27.02	33.57	-0.36	27.13
44.25	-0.30	27.02	40.07	-0.67	27.11	37.04	-0.49	27.01	34.56	-0.37	27.13
45.31	-0.25	27.03	41.02	-0.70	27.13	38.94	-0.54	27.04	35.54	-0.38	27.14
46.13	-0.25	27.03	42.42	-0.71	27.14	39.88	-0.58	27.06	36.55	-0.39	27.14
47.94	-0.26	27.05	43.53	-0.72	27.14	40.65	-0.61	27.07	37.07	-0.38	27.14
48.05	-0.27	27.06	44.59	-0.73	27.15	41.15	-0.63	27.08	38.37	-0.39	27.15
49.25	-0.27	27.05				42.05	-0.74	27.16	39.38	-0.40	27.15
50.12	-0.26	27.04							40.32	-0.40	27.14
									41.34	-0.43	27.16
									42.34	-0.44	27.16
									43.27	-0.45	27.17
									45.38	-0.45	27.17
									46.17	-0.50	27.17

Appendix 3.1-1. Temperature and Salinity Data for the AEMP Expansion Sites, Doris North Project, 2011

RB2											
23-Apr-11			21-Jul-11			14-Aug-11			22-Sep-11		
Depth (m)	Temperature (°C)	Salinity	Depth (m)	Temperature (°C)	Salinity	Depth (m)	Temperature (°C)	Salinity	Depth (m)	Temperature (°C)	Salinity
1.56	-1.41	26.84	0.93	10.80	19.99	0.98	7.59	24.44	0.61	6.18	23.97
1.86	-1.41	26.84	1.57	6.61	23.87	1.51	9.69	23.15	1.03	3.59	25.83
2.03	-1.41	26.84	2.14	4.84	24.95	2.12	9.73	23.10	1.49	3.58	25.85
2.89	-1.41	26.84	2.48	4.29	25.39	2.50	9.58	23.23	2.22	3.58	25.85
3.29	-1.41	26.84	3.02	3.34	25.86	3.20	9.07	23.95	2.78	3.57	25.85
3.97	-1.41	26.84	3.44	2.96	26.11	3.61	8.79	24.37	3.16	3.56	25.85
4.01	-1.41	26.84	4.00	2.50	26.32	4.02	8.43	24.44	3.38	3.56	25.85
4.81	-1.41	26.84	4.72	2.42	26.35	4.78	7.57	24.50	4.10	3.55	25.85
5.34	-1.41	26.84	5.01	2.35	26.32	5.23	6.58	24.29	4.71	3.51	25.86
5.84	-1.41	26.84	5.29	2.08	26.31	5.66	5.14	25.16	5.31	3.46	25.89
6.73	-1.41	26.85	5.86	1.43	26.11	6.09	4.32	25.63	5.88	3.32	25.94
7.06	-1.41	26.84	6.01	1.05	26.30	6.59	3.79	25.61	6.28	3.31	25.94
7.59	-1.41	26.84	6.63	0.40	26.68	7.21	3.04	25.99	6.26	3.30	25.95
8.65	-1.41	26.84	7.38	0.39	26.68	7.74	2.79	26.16	6.87	3.25	25.97
9.10	-1.41	26.84	7.79	0.33	26.73	8.18	2.63	26.41	7.28	3.16	26.03
9.64	-1.41	26.84	8.07	0.33	26.74	8.72	2.55	26.24	7.74	3.11	26.06
10.98	-1.41	26.84	8.31	0.32	26.74	9.21	2.24	26.28	8.18	3.06	26.09
11.70	-1.40	26.84	9.22	0.13	26.73	9.70	2.01	26.29	8.60	3.02	26.12
13.26	-1.40	26.84	9.72	0.16	26.84	10.06	1.72	26.42	9.04	3.01	26.13
14.23	-1.40	26.84	10.18	0.19	26.81	11.25	1.43	26.54	9.79	3.01	26.14
15.48	-1.40	26.84	11.12	0.08	26.79	12.13	1.12	26.58	10.03	3.00	26.14
16.59	-1.40	26.84	13.23	-0.07	26.85	13.11	0.81	26.66	11.17	2.96	26.16
17.48	-1.39	26.84	14.49	-0.21	26.79	14.06	0.62	26.69	12.13	2.93	26.16
18.06	-1.39	26.83	15.26	-0.45	26.86	15.20	0.38	26.79	13.03	2.92	26.25
19.22	-1.40	26.84	16.36	-0.59	26.86	16.47	0.21	26.79	14.03	2.95	26.25
20.25	-1.39	26.83	17.10	-0.67	26.90	17.15	0.10	26.83	15.01	2.91	26.25
21.72	-1.39	26.83	18.14	-0.69	26.90	18.15	-0.03	26.89	16.06	2.69	26.30
23.06	-1.38	26.82	19.04	-0.67	26.95	19.17	-0.06	26.90	17.30	2.61	26.34
24.40	-1.38	26.84	20.29	-0.59	27.02	20.14	-0.09	26.93	18.53	2.54	26.31
25.53	-1.38	26.82	21.13	-0.38	27.02	21.10	-0.13	26.92	19.33	2.44	26.29
26.61	-1.36	26.82	22.20	-0.37	27.00	22.17	-0.21	26.92	20.48	2.22	26.33
27.96	-1.33	26.81	23.53	-0.38	27.02	23.55	-0.28	26.96	21.24	2.00	26.39
29.24	-1.27	26.84	24.21	-0.35	27.02	24.09	-0.30	26.96	22.27	1.57	26.61
29.86	-0.62	26.89	25.02	-0.34	27.01	25.15	-0.34	26.99	22.21	1.49	26.65
31.51	-0.61	26.89	26.84	-0.37	27.03	26.32	-0.40	27.01	22.67	1.39	26.52
32.97	-0.49	26.90	27.86	-0.36	27.04	27.19	-0.31	27.08	23.24	1.11	26.68
34.15	-0.47	26.92	28.79	-0.37	27.02	28.43	-0.28	27.04	24.41	0.97	26.72
35.12	-0.29	26.83	30.11	-0.40	27.03	29.44	-0.30	27.04	25.24	0.77	26.80
36.59	-0.26	26.95	31.65	-0.43	27.04	30.05	-0.32	27.03	26.20	0.58	26.94
37.95	-0.25	26.96	33.43	-0.45	27.06	31.11	-0.37	27.05	27.46	0.31	26.85
39.16	-0.25	26.94	35.44	-0.47	27.06	32.29	-0.40	27.06	28.22	0.11	26.93
40.81	-0.22		37.14	-0.51	27.07	33.31	-0.48	27.05	29.65	-0.07	26.97
42.31	-0.22		38.49	-0.57	27.07	34.01	-0.54	27.06	30.16	-0.17	27.06
43.81	-0.23		40.39	-0.63	27.07	35.30	-0.63	27.08	31.82	-0.34	27.11
44.07	-0.22		41.25	-0.67	27.09	36.52	-0.70	27.10	32.86	-0.39	27.09
45.21	-0.24		43.35	-0.71	27.11	37.03	-0.72	27.11	34.09	-0.45	27.13
46.45	-0.25		45.03	-0.74	27.12	38.39	-0.75	27.13	35.47	-0.51	27.14
47.78	-0.26		46.82	-0.74	27.12	39.54	-0.77	27.14	36.47	-0.60	27.21
48.98	-0.24		47.57	-0.75	27.14	40.04	-0.79	27.14	38.08	-0.63	27.20
49.10	-0.23		48.17	-0.75	27.14	41.44	-0.80	27.16	39.19	-0.66	27.22
50.26	-0.31		49.96	-0.77	27.14	41.93	-0.80	27.17	40.17	-0.68	27.24
51.53	-0.33		50.50	-0.78	27.15	42.16	-0.81	27.17	41.97	-0.70	27.26
52.91	-0.34		51.65	-0.77	27.16	43.49	-0.83	27.18	42.31	-0.70	27.27
53.02	-0.33		52.33	-0.78	27.16	44.06	-0.84	27.18	43.08	-0.70	27.27
54.17	-0.26		53.03	-0.78	27.16	45.21	-0.84	27.19	44.50	-0.72	27.28
55.03	-0.23		54.09	-0.78	27.16	46.29	-0.86	27.19	45.20	-0.73	27.30
56.46	-0.26		55.34	-0.78	27.16	47.02	-0.87	27.19	46.30	-0.73	27.30
57.41	-0.24		56.55	-0.79	27.16	48.23	-0.88	27.19	47.40	-0.72	27.33
58.04	-0.30		57.18	-0.80	27.16	49.35	-0.89	27.20	48.11	-0.71	27.33
59.30	-0.26		58.48	-0.80	27.17	50.09	-0.90	27.20	49.21	-0.71	27.34
			59.57	-0.80	27.17	51.33	-0.92	27.21	50.58	-0.70	27.35
			60.40	-0.80	27.17	52.24	-0.92	27.21	51.13	-0.70	27.35
			61.95	-0.81	27.17	53.13	-0.94	27.22	52.14	-0.71	27.37
			63.15	-0.83	27.18	54.34	-0.95	27.22	53.19	-0.70	27.37
			64.26	-0.83	27.18	55.67	-0.95	27.22	54.59	-0.69	27.38
			65.79	-0.84	27.18	56.44	-0.96	27.23	55.26	-0.68	27.38
			66.56	-0.83	27.19	57.24	-0.96	27.24	56.50	-0.68	27.37
			67.49	-0.83	27.19	58.68	-0.96	27.24	57.55	-0.69	27.38
			68.94	-0.84	27.19	59.45	-0.96	27.24	58.18	-0.68	27.39
			69.60	-0.85	27.20	60.11	-0.96	27.24	59.00	-0.67	27.39
			70.42	-0.85	27.20	61.23	-0.96	27.25	60.62	-0.67	27.39
			71.22	-0.86	27.20	62.10	-0.96	27.25	61.82	-0.67	27.39
						63.46	-0.96	27.26	62.50	-0.67	27.40
									63.03	-0.67	27.41
									64.24	-0.67	27.41
									65.03	-0.67	27.42
									66.50	-0.65	27.44
									67.21	-0.64	27.44
									68.34	-0.62	27.44
									69.15	-0.61	27.44

Appendix 3.1-1. Temperature and Salinity Data for the AEMP Expansion Sites, Doris North Project, 2011

REF-Marine 2											
April			21-Jul-11			19-Aug-11			25-Sep-11		
Depth	Temperature	Salinity	Depth	Temperature	Salinity	Depth	Temperature	Salinity	Depth	Temperature	Salinity
(m)	(°C)		(m)	(°C)		(m)	(°C)		(m)	(°C)	
Data not collected.			0.94	9.58	15.67	0.80	4.54	24.51	0.88	2.38	26.40
			1.49	5.97	23.53	1.09	2.74	25.51	1.04	3.19	25.77
			2.26	4.51	24.58	1.76	1.47	26.28	1.45	3.34	25.64
			2.89	3.07	25.15	2.28	1.27	26.49	2.00	3.34	25.63
			3.00	2.49	25.71	2.68	1.15	26.55	2.74	3.33	25.66
			3.64	1.80	26.04	3.09	1.04	26.62	3.30	3.33	25.71
			4.04	1.66	26.14	3.45	0.98	26.67	3.73	3.31	25.81
			4.49	1.47	26.21	4.35	0.85	26.63	4.38	3.25	25.87
			5.21	1.60	26.50	4.61	0.73	26.72	4.58	3.24	25.88
			5.65	1.59	26.54	4.99	0.63	26.72	5.06	3.24	25.88
			6.13	1.37	26.38	5.50	0.55	26.76	5.44	3.24	25.88
			6.48	1.09	26.58	6.13	0.50	26.76	6.15	3.32	25.78
			7.08	0.78	26.61	6.68	0.45	26.80	6.50	3.24	25.88
			7.53	0.61	26.71	7.16	0.42	26.82	7.15	3.24	25.89
			8.01	0.50	26.65	7.79	0.40	26.83	7.53	3.23	25.89
			8.63	0.28	26.79	8.28	0.39	26.85	8.30	3.23	25.90
			9.17	0.21	26.89	9.11	0.43	26.90	9.32	3.24	25.93
			9.68	0.05	26.84	9.88	0.44	26.87	10.04	3.25	25.93
			10.02	0.04	26.84	10.42	0.43	26.88	10.56	3.26	25.93
			11.36	-0.30	26.77	11.27	0.42	26.91	11.22	3.26	25.93
			12.33	-0.52	26.82	12.14	0.44	26.90	12.09	3.26	25.93
			13.37	-0.59	26.85	13.59	0.40	26.89	13.17	3.26	25.93
			14.36	-0.67	26.74	14.47	0.33	26.87	14.09	3.26	25.94
			15.15	-0.82	26.83	15.04	0.26	26.89	15.01	3.26	25.93
			16.25	-0.97	26.89	16.72	0.19	26.90	16.32	3.27	25.94
			17.27	-0.98	26.93	17.24	0.17	26.92	17.54	3.27	25.95
			18.09	-0.93	26.92	18.07	0.13	26.92	18.43	3.27	25.94
			19.09	-0.84	26.96	19.30	0.09	26.93	19.15	3.27	25.95
			20.06	-0.81	26.99	20.67	0.04	26.93	20.07	3.26	25.95
			21.06	-0.80	26.96	21.35	-0.03	26.89	20.65	3.24	25.97
			22.16	-0.79	26.97	22.03	-0.12	26.95	22.04	3.23	25.97
			23.06	-0.79	26.99	23.11	-0.20	26.97	22.85	3.19	25.99
			24.03	-0.77	26.95	24.28	-0.25	26.95	24.07	3.16	25.99
			25.01	-0.74	27.00	25.01	-0.29	26.97	25.05	3.09	26.03
			26.11	-0.71	26.99	26.66	-0.39	26.97	26.89	2.10	25.90
			27.22	-0.69	27.00	27.42	-0.43	27.01	28.07	1.09	26.46
			28.07	-0.69	26.98	28.62	-0.46	27.03	29.76	0.34	26.82
			29.91	-0.68	27.01	29.18	-0.47	27.04	30.05	-0.04	27.09
			30.63	-0.68	27.01	30.51	-0.48	27.04	31.39	-0.07	27.07
			31.06	-0.66	27.04	31.20	-0.49	27.04	32.48	-0.16	27.07
			32.15	-0.60	27.03	32.79	-0.49	27.04	33.04	-0.25	27.12
			33.21	-0.58	27.05	33.55	-0.49	27.03	34.48	-0.36	27.17
			34.10	-0.56	27.05	34.29	-0.50	27.05	35.30	-0.36	27.15
			35.07	-0.54	27.05	35.07	-0.50	27.04	36.98	-0.40	27.16
			36.21	-0.52	27.06	36.56	-0.51	27.10	37.93	-0.43	27.17
			37.33	-0.48	27.08	37.31	-0.50	27.07	38.69	-0.44	27.18
		38.65	-0.47	27.08	38.10	-0.49	27.07	39.01	-0.46	27.19	
		39.49	-0.45	27.10	39.74	-0.49	27.08	40.24	-0.47	27.20	
		40.17	-0.42	27.12	40.45	-0.49	27.06	41.50	-0.47	27.20	
		41.69	-0.39	27.10	41.02	-0.51	27.08				
		42.44	-0.39	27.11	42.67	-0.56	27.07				
		43.18	-0.35	27.14	43.03	-0.59	27.10				

Appendix 3.1-2

**Dissolved Oxygen Concentrations and Percent Saturation
for the Marine AEMP Expansion Sites, Doris North
Project, 2011**

Appendix 3.1-2. Dissolved Oxygen Concentrations and Percent Saturation for the AEMP Expansion Sites, Doris North Project, 2011

RB1											
23-Apr-11			21-Jul-11			14-Aug-11			22-Sep-11		
Depth	Dissolved Oxygen	Dissolved Oxygen Saturation	Depth	Dissolved Oxygen	Dissolved Oxygen Saturation	Depth	Dissolved Oxygen	Dissolved Oxygen Saturation	Depth	Dissolved Oxygen	Dissolved Oxygen Saturation
(m)	(mg/L)	(%)	(m)	(mg/L)	(%)	(m)	(mg/L)	(%)	(m)	(mg/L)	(%)
1.50	11.90	94.5	1.05	9.87	96.7	0	9.01	85.5	0	11.23	85.3
1.88	11.90	94.5	1.76	10.71	96.3	1	9.03	85.3	0.5	11.46	86.7
2.33	11.89	94.4	2.12	11.48	103.8	2	9.06	85.2	1	11.48	86.8
2.92	11.89	94.4	2.72	11.98	107.2	3	9.08	85.0	1.5	11.49	86.9
3.88	11.90	94.5	3.13	12.13	108.7	4	9.21	84.5	2	11.54	86.8
4.81	11.88	94.3	3.60	12.23	108.2	5	9.29	83.2	2.5	11.61	86.9
5.65	11.88	94.3	4.24	12.24	107.2	6	9.41	83.2	3	11.62	87
5.92	11.88	94.4	5.01	12.34	106.6	7	9.44	83.2	3.5	11.63	87.1
6.80	11.88	94.3	5.53	12.27	104.2	8	9.43	82.7	4	11.67	87.1
7.77	11.89	94.4	6.04	12.27	103.5	9	9.46	82.8	4.5	11.67	87.2
8.80	11.88	94.3	6.68	12.42	105.0	10	9.50	82.8	5	11.67	86.9
9.50	11.87	94.3	7.10	12.50	105.7	11	9.49	82.8	5.5	11.71	86.9
10.14	11.87	94.3	7.52	12.48	105.0	12	9.49	82.8	6	11.71	87
10.81	11.86	94.2	8.15	12.45	104.7	13	9.50	82.7	6.5	11.71	86.9
11.70	11.86	94.2	8.85	12.48	104.8	14	9.50	82.5	7	11.70	86.9
11.92	11.86	94.2	9.32	12.47	104.8	15	9.51	82.4	7.5	11.69	86.8
13.12	11.85	94.1	9.78	12.50	105.0	16	9.51	82.4	8	11.69	86.8
14.25	11.85	94.1	10.20	12.51	105.2	17	9.53	82.3	8.5	11.68	86.8
15.78	11.84	94.0	11.17	12.39	103.2	18	9.52	82.0	9	11.71	86.7
16.40	11.82	93.9	13.23	12.29	101.6	19	9.48	81.2	9.5	11.70	86.7
17.63	11.83	93.9	14.06	12.24	100.9	20	9.44	80.4	10	11.73	86.6
18.51	11.78	93.6	15.07	12.19	100.3	22	9.31	78.5	11	11.73	86.7
19.41	11.82	94.0	16.24	12.08	99.3	24	9.16	76.9	12	11.71	86.6
20.09	11.84	94.0	17.09	12.04	98.8	26	9.08	76.2	13	11.72	86.6
21.14	11.81	93.8	18.17	11.94	97.8	28	9.07	75.9	14	11.71	86.5
22.29	11.80	93.7	19.59	11.86	96.8	30	8.99	74.8	15	11.64	85.8
23.43	11.78	93.6	20.12	11.84	96.6	32	8.87	73.2	20	11.41	80.4
24.51	11.79	93.7	21.45	11.56	94.2	34	8.77	72.7	25	11.11	76.4
25.72	11.58	91.9	22.43	11.48	93.6	36	8.64	71	30	10.80	73.6
27.22	11.53	91.6	23.29	11.45	93.3				35	10.65	72.2
28.31	11.47	91.1	24.22	11.29	92.2				40	10.52	71.2
29.26	10.67	84.9	25.36	11.07	90.4				45	10.42	70.3
30.29	10.48	83.6	26.59	10.79	88.1						
31.56	10.26	82.4	27.47	10.63	86.9						
32.78	10.09	82.1	28.11	10.49	85.7						
33.91	10.16	83.1	29.45	10.25	83.7						
34.48	10.10	82.7	30.49	10.06	82.1						
35.19	10.09	82.6	31.39	9.90	80.9						
36.66	10.07	82.4	32.08	9.83	80.3						
37.17	10.06	82.3	33.27	9.61	78.4						
38.03	10.05	82.3	34.36	9.55	78.0						
39.15	10.04	82.2	35.22	9.51	77.6						
40.73	10.01	81.9	36.30	9.39	76.4						
41.86	9.99	81.8	37.76	9.33	75.9						
42.83	9.97	81.6	38.76	9.31	75.7						
44.25	9.93	81.3	40.07	9.32	75.6						
45.31	9.88	81.0	41.02	9.32	75.6						
46.13	9.87	81.0	42.42	9.34	75.7						
47.94	9.87	81.0	43.53	9.35	75.8						
48.05	9.87	80.9	44.59	9.35	75.8						
49.25	9.88	81.0									
50.12	9.92	81.3									

Note: April and July DO data comes from CTD probe, August and September DO data are from YSI probe.
All data are below marine CCME guidelines for aquatic life (8.0 mg/L).

Appendix 3.1-2. Dissolved Oxygen Concentrations and Percent Saturation for the AEMP Expansion Sites, Doris North Project, 2011

RB2											
23-Apr-11			21-Jul-11			14-Aug-11			22-Sep-11		
Depth	Dissolved Oxygen	Dissolved Oxygen Saturation	Depth	Dissolved Oxygen	Dissolved Oxygen Saturation	Depth	Dissolved Oxygen	Dissolved Oxygen Saturation	Depth	Dissolved Oxygen	Dissolved Oxygen Saturation
(m)	(mg/L)	(%)	(m)	(mg/L)	(%)	(m)	(mg/L)	(%)	(m)	(mg/L)	(%)
1.56	11.90	94.4	0.93	9.07	93.1	0	8.93	85.4	0	11.44	86.5
1.86	11.89	94.4	1.57	9.22	88.0	1	8.95	85.2	0.5	11.49	86.9
2.03	11.89	94.4	2.14	9.20	84.8	2	9.05	83.8	1	11.51	87.1
2.89	11.89	94.4	2.48	9.70	88.4	3	9.16	84.2	1.5	11.56	87.2
3.29	11.88	94.3	3.02	10.24	91.5	4	9.20	83.4	2	11.57	87.3
3.97	11.89	94.3	3.44	10.74	95.2	5	9.27	83.5	2.5	11.57	87.3
4.01	11.88	94.3	4.00	11.09	97.2	6	9.39	82.0	3	11.57	87.3
4.81	11.89	94.3	4.72	11.50	100.7	7	9.44	81.9	3.5	11.57	87.4
5.34	11.88	94.3	5.01	11.57	101.1	8	9.46	82.0	4	11.61	87.4
5.84	11.88	94.3	5.29	11.64	101.0	9	9.49	82.0	4.5	11.61	87.2
6.73	11.88	94.3	5.86	11.68	99.5	10	9.49	82.0	5	11.62	87.3
7.06	11.88	94.3	6.01	11.68	98.6	11	9.50	81.9	5.5	11.65	87.3
7.59	11.88	94.3	6.63	11.96	99.5	12	9.48	81.4	6	11.69	87.3
8.65	11.87	94.2	7.38	12.14	101.1	13	9.48	81.0	6.5	11.69	87.3
9.10	11.87	94.2	7.79	12.25	101.9	14	9.45	80.4	7	11.72	87.3
9.64	11.87	94.2	8.07	12.27	102.0	15	9.44	80.0	7.5	11.72	87.3
10.98	11.86	94.1	8.31	12.32	102.4	16	9.40	79.2	8	11.72	87.3
11.70	11.87	94.2	9.22	12.28	101.5	17	9.39	78.8	8.5	11.72	87.3
13.26	11.86	94.2	9.72	12.35	102.3	18	9.40	78.6	9	11.74	87.2
14.23	11.86	94.1	10.18	12.36	102.4	19	9.34	78.0	9.5	11.73	87.2
15.48	11.85	94.1	11.12	12.29	101.5	20	9.33	77.8	10	11.72	87.1
16.59	11.84	94.0	13.23	12.21	100.5	22	9.24	76.7	11	11.72	87.1
17.48	11.84	94.0	14.49	12.20	100.0	24	9.18	76.2	12	11.73	87.1
18.06	11.82	93.9	15.26	12.13	98.9	26	9.11	75.6	13	11.74	87.2
19.22	11.76	93.3	16.36	12.07	98.0	28	9.06	75.2	14	11.72	87.1
20.25	11.74	93.2	17.10	11.84	96.0	30	9.04	75.0	15	11.72	86.4
21.72	11.72	93.1	18.14	11.79	95.5	32	9.03	74.8	20	11.54	82.8
23.06	11.70	92.9	19.04	11.80	95.6	34	8.99	74.3	25	11.27	78.6
24.40	11.56	91.8	20.29	11.72	95.2	36	8.92	73.7	30	10.91	74.5
25.53	11.45	90.9	21.13	11.50	94.0	38	8.83	72.9	35	10.66	72
26.61	11.38	90.5	22.20	11.12	90.9	40	8.70	71.4	40	10.39	70
27.96	11.29	89.8	23.53	10.78	88.1	44	8.64	70.7	45	10.34	69.3
29.24	11.21	89.3	24.21	10.63	86.9	48	8.42	67.9	50	10.36	69.5
29.86	10.42	84.5	25.02	10.18	83.3	52	8.27	66.7	55	10.34	69.3
31.51	10.37	84.2	26.84	9.60	78.5	56	8.24	66.5	60	10.36	69.5
32.97	10.32	84.0	27.86	9.58	78.3	60	8.24	66.5	65	10.35	69.6
34.15	10.33	84.1	28.79	9.52	77.9	62	8.25	66.5	70	10.36	69.5
35.12	10.10	82.6	30.11	9.48	77.4				75	10.15	68.2
36.59	10.09	82.7	31.65	9.43	77.0						
37.95	10.08	82.6	33.43	9.41	76.8						
39.16	10.06	82.4	35.44	9.36	76.3						
40.81	10.01	74.2	37.14	9.31	75.9						
42.31	10.00	74.1	38.49	9.24	75.2						
43.81	9.99	73.9	40.39	9.23	74.9						
44.07	9.94	73.2	41.25	9.25	75.0						
45.21	9.93	73.1	43.35	9.28	75.2						
46.45	9.92	73.0	45.03	9.37	75.9						
47.78	9.91	72.8	46.82	9.41	76.2						
48.98	9.87	72.4	47.57	9.46	76.6						
49.10	9.87	72.4	48.17	9.48	76.8						
50.26	9.87	72.2	49.96	9.50	76.9						
51.53	9.84	71.9	50.50	9.54	77.2						
52.91	9.83	71.8	51.65	9.58	77.5						
53.02	9.79	71.5	52.33	9.60	77.7						
54.17	9.69	70.8	53.03	9.63	77.9						
55.03	9.66	70.6	54.09	9.64	78.0						
56.46	9.63	70.3	55.34	9.66	78.2						
57.41	9.41	68.6	56.55	9.68	78.3						
58.04	9.56	69.7	57.18	9.70	78.5						
59.30	9.34	68.1	58.48	9.72	78.6						
60.01	9.29	67.7	59.57	9.74	78.8						
61.52	9.21	66.9	60.40	9.75	78.8						
62.70	9.21	65.9	61.95	9.75	78.9						
63.01	9.23	65.7	63.15	9.77	79.0						
63.01	9.32	66.1	64.26	9.77	79.0						
64.61	9.36	66.3	65.79	9.79	79.1						
65.83	9.36	66.3	66.56	9.80	79.2						
66.00	9.37	66.3	67.49	9.79	79.2						
			68.94	9.80	79.2						
			69.60	9.80	79.2						
			70.42	9.81	79.2						
			71.22	9.82	79.3						

Appendix 3.1-2. Dissolved Oxygen Concentrations and Percent Saturation for the AEMP Expansion Sites, Doris North Project, 2011

REF-Marine 2											
April			21-Jul-11			19-Aug-11			Sept		
Depth	Dissolved Oxygen	Dissolved Oxygen Saturation	Depth	Dissolved Oxygen	Dissolved Oxygen Saturation	Depth	Dissolved Oxygen	Dissolved Oxygen Saturation	Depth	Dissolved Oxygen	Dissolved Oxygen Saturation
(m)	(mg/L)	(%)	(m)	(mg/L)	(%)	(m)	(mg/L)	(%)	(m)	(mg/L)	(%)
Data not collected.			0.94	10.89	105.8	0.0	11.57	79.1	Data not collected.		
			1.49	10.61	99.5	1.0	9.36	78.4			
			2.26	10.25	93.4	2.0	9.36	78.8			
	2.89	10.53	92.9	3.0	9.37	78.4					
	3.00	10.86	94.9	3.9	9.30	77.6					
	3.64	11.23	96.6	4.9	9.23	76.9					
	4.04	11.51	98.7	5.9	9.18	76.5					
	4.49	11.55	98.6	6.9	9.18	76.3					
	5.21	11.76	100.9	7.9	9.15	76.3					
	5.65	11.92	102.3	8.9	9.15	76.3					
	6.13	11.92	101.6	9.8	9.16	76.4					
	6.48	12.01	101.8	10.8	9.20	76.7					
	7.08	11.93	100.3	11.8	9.18	76.4					
	7.53	12.01	100.6	12.8	9.11	75.8					
	8.01	12.20	101.8	13.8	9.10	75.5					
	8.63	12.14	100.8	14.8	9.09	75.2					
	9.17	12.21	101.3	15.8	9.06	75.0					
	9.68	12.19	100.7	16.7	9.04	74.8					
	10.02	12.10	99.9	17.7	9.01	74.6					
	11.36	11.99	98.0	18.7	8.99	74.2					
	12.33	11.98	97.4	19.7	8.90	73.5					
	13.37	12.03	97.6	20.7	8.82	72.6					
	14.36	11.93	96.5	23.6	8.50	69.6					
	15.15	11.95	96.4	25.6	7.76	63.2					
	16.25	11.93	95.8	27.6	7.50	61.1					
	17.27	11.88	95.5	29.5	7.00	56.9					
	18.09	11.74	94.5	31.5	6.79	55.2					
	19.09	11.61	93.7	33.5	6.76	55.0					
	20.06	11.44	92.4	35.5	6.70	54.5					
	21.06	11.22	90.6	37.4	6.76	55.0					
	22.16	10.98	88.7	39.4	6.79	55.2					
	23.06	10.82	87.4	41.4	6.63	53.9					
	24.03	10.68	86.3								
	25.01	10.52	85.1								
	26.11	10.36	83.9								
	27.22	10.15	82.2								
	28.07	10.06	81.5								
	29.91	9.96	80.7								
	30.63	9.90	80.3								
	31.06	9.88	80.1								
	32.15	9.74	79.1								
	33.21	9.58	77.9								
	34.10	9.44	76.8								
	35.07	9.31	75.8								
	36.21	9.16	74.6								
	37.33	9.01	73.5								
38.65	8.83	72.0									
39.49	8.74	71.3									
40.17	8.63	70.5									
41.69	8.36	68.4									
42.44	8.22	67.2									
43.18	8.07	66.1									

Appendix 3.2-1

**Water Quality Data for Marine AEMP Expansion Sites,
Doris North Project, 2011**

Appendix 3.2-1. Water Quality Data for Marine AEMP Expansion Sites, 2011

Site ID:				RB1	RB1	RB1	RB1	RB1	RB2	RB2	RB2	RB2	RB2	RB1	RB1	RB1	RB1	RB1	RB1	RB2	RB2	RB2	RB2	REF-Marine 2	REF-Marine 2	REF-Marine 2
Depth Zone:				Surface	Surface	Above Pycnocline	Below Pycnocline	Deep	Surface	Above Pycnocline	Above Pycnocline	Below Pycnocline	Deep	Surface	Surface	Above Pycnocline	Below Pycnocline	Deep	Deep	Surface	Above Pycnocline	Below Pycnocline	Deep	Surface	Above Pycnocline	Below Pycnocline
Depth Sampled (m):				3	3	25	33	48	3	35	35	44	48	1	1	7	15	40	40	1	7	15	40	1	8	20
Replicate:				1	2	1	1	1	1	1	2	1	1	1	2	1	1	1	2	1	1	1	1	1	1	1
Date Sampled:		CCME Guideline for the Protection of Aquatic Life ^a	Realized Detection Limit	24-Apr-11	24-Apr-11	24-Apr-11	24-Apr-11	24-Apr-11	23-Apr-11	23-Apr-11	23-Apr-11	23-Apr-11	23-Apr-11	22-Jul-11	22-Jul-11	22-Jul-11	22-Jul-11	22-Jul-11	22-Jul-11	22-Jul-11	22-Jul-11	22-Jul-11	22-Jul-11	27-Jul-11	27-Jul-11	27-Jul-11
ALS Sample ID:	Units			L998223-12	L998223-13	L998223-11	L998223-10	L998223-9	L998223-17	L998223-5	L998223-6	L998223-7	L998223-8	L1035865-6	L1035865-5	L1035865-4	L1035865-3	L1035865-2	L1035865-1	L1035865-10	L1035865-9	L1035865-8	L1035865-7	L1037843-1	L1037843-2	L1037843-3
Physical Tests																										
Hardness (as CaCO ₃)	mg/L		4.3	4850	4810	4760	4990	4800	4880	4850	4830	4860	4890	2830	2760	4230	4890	5090	5070	2890	3730	4900	5100	3000	3210	4440
pH	pH	7.0-8.7	0.10	7.79	7.80	7.79	7.75	7.76	7.81	7.79	7.77	7.77	7.77	7.82	7.83	7.86	7.88	7.70	7.82	7.84	7.87	7.88	7.81	8.04	8.00	7.83
Salinity (EC)	g/L	<10% fluctuation ^b	1.0	26.1	26.0	26.0	26.1	26.3	26.1	26.0	26.2	26.5	26.3	15.3	14.9	23.2	26.4	27.0	27.0	15.9	20.0	26.5	27.2	16.0	18.0	24.9
Total Suspended Solids	mg/L	dependent on background levels	2.0 to 4.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	2.2	<2.0	2.0	2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0
Turbidity	NTU	dependent on background levels	0.10	0.27	0.20	0.18	0.26	0.21	0.16	0.27	0.15	0.31	0.23	0.41	0.43	0.60	0.49	0.40	0.35	0.49	0.42	0.41	0.22	0.54	0.48	0.50
Anions and Nutrients																										
Alkalinity, Total (as CaCO ₃)	mg/L		2.0	127	125	130	132	136	134	132	134	129	133	47.0	46.6	104	116	119	118	47.9	95.6	117	118	48.5	86.6	110
Ammonia (as N)	mg/L		0.0050	0.0062	<0.0050	<0.0050	<0.0050	<0.0050	<0.0050	<0.0050	<0.0050	<0.0050	<0.0050	<0.0050	<0.0050	<0.0050	<0.0050	<0.0050	<0.0050	<0.0050	<0.0050	<0.0050	0.0261	0.0256	0.0302	
Bromide (Br)	mg/L		5.0	45.0	42.0	42.4	43.0	45.1	41.8	43.3	43.3	42.5	41.9	26.9	26.2	41.2	48.2	50.8	51.4	28.2	38.6	49.8	48.9	36.8	42.1	56.4
Chloride (Cl)	mg/L		50	15300	15400	15300	15200	15500	15200	15500	15600	15800	15500	8470	8140	12700	14700	15000	14900	8950	11600	15100	15100	10100	11600	15800
Fluoride (F)	mg/L	0.03 or 0.75		0.89	0.90	0.86	0.94	0.90	0.92	0.98	0.84	0.89	0.90	<0.75	<0.75	0.85	1.15	0.82	0.81	<0.75	<0.75	0.86	0.84	0.760	0.670	0.860
Nitrate and Nitrite (as N)	mg/L		0.0060	0.0409	0.0402	0.0396	0.0575	0.0749	0.0390	0.0688	0.0654	0.0608	0.0674	<0.0060	<0.0060	<0.0060	<0.0060	0.0802	0.0793	<0.0060	<0.0060	<0.0060	0.0786	<0.0060	<0.0060	<0.0060
Nitrate (as N)	mg/L	3.612 ^b	0.0060	0.0384	0.0402	0.0373	0.0575	0.0749	0.0390	0.0688	0.0654	0.0608	0.0674	<0.0060	<0.0060	<0.0060	<0.0060	0.0802	0.0793	<0.0060	<0.0060	<0.0060	0.0786	<0.0060	<0.0060	<0.0060
Nitrite (as N)	mg/L		0.0020	0.0025	<0.0020	0.0023	<0.0020	<0.0020	<0.0020	<0.0020	<0.0020	<0.0020	<0.0020	<0.0020	<0.0020	<0.0020	<0.0020	<0.0020	<0.0020	<0.0020	<0.0020	<0.0020	<0.0020	<0.0020	<0.0020	<0.0020
Total Phosphorus	mg/L		0.0020	0.0378	0.0374	0.0375	0.0413	0.0438	0.0383	0.0453	0.0451	0.0453	0.0468	0.0202	0.0207	0.0325	0.0357	0.0464	0.0485	0.0205	0.0276	0.0371	0.0463	0.0221	0.0227	0.0385
Sulphate (SO ₄)	mg/L		50	2110	2110	2100	2100	2130	2090	2130	2140	2180	2130	1190	1140	1810	2090	2130	2130	1270	1650	2150	2120	1350	1560	2140
Cyanides																										
Total Cyanide	mg/L		0.0010	0.0013	0.0010	0.0013	0.0013	0.0012	0.0013	0.0012	0.0014	0.0013	0.0012	0.0013	0.0016	0.0016	0.0014	<0.0010	<0.0010	0.0021	0.0016	0.0013	<0.0010	0.0019	0.0018	0.0013
Free Cyanide	mg/L		0.0010 or 0.0050	<0.0050	<0.0050	<0.0050	<0.0050	<0.0050	<0.0050	<0.0050	<0.0050	<0.005	<0.0050	<0.0050	<0.0050	<0.0050	<0.0050	<0.0050	<0.0050	<0.0050	<0.0050	<0.0050	<0.0050	<0.0050	<0.0050	<0.0050
Organic / Inorganic Carbon																										
Total Organic Carbon	mg/L		0.50	1.14	1.15	1.09	1.30	1.15	1.16	1.15	1.17	1.17	1.12	1.76	1.77	1.56	1.46	1.39	1.39	1.77	1.67	1.73	1.42	1.81	1.65	1.50
Total Metals																										
Aluminum (Al)	mg/L		0.0050	<0.0050	<0.0050	<0.0050	<0.0050	<0.0050	<0.0050	<0.0050	<0.0050	<0.0050	<0.0050	0.0077	0.0086	0.0080	0.0090	0.0068	0.0093	0.0086	0.0077	0.0066	<0.0050	0.0135	0.0072	<0.0050
Antimony (Sb)	mg/L		0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050
Arsenic (As)	mg/L	0.0125 ^b	0.0004	0.00101	0.00107	0.00102	0.00094	0.00094	0.00104	0.00106	0.00100	0.00119	0.00100	0.00068	0.00072	0.00103	0.00115	0.00123	0.00122	0.00070	0.00088	0.00115	0.00116	0.00079	0.00095	0.00110
Barium (Ba)	mg/L		0.0010	0.0112	0.0111	0.0113	0.0116	0.0133	0.0118	0.0117	0.0122	0.0129	0.0129	0.0078	0.0077	0.0109	0.0121	0.0127	0.0127	0.0075	0.0094	0.0118	0.0129	0.0080	0.0091	0.0111
Beryllium (Be)	mg/L		0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050
Bismuth (Bi)	mg/L		0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050
Boron (B)	mg/L		0.10	3.78	3.77	3.82	3.64	3.83	3.92	3.56	3.76	3.78	3.76	2.16	2.17	3.32	3.72	3.82	3.84	2.18	2.81	3.83	3.99	2.25	2.60	3.47
Cadmium (Cd)	mg/L	0.00012	0.000020	0.000045	0.000049	0.000047	0.000049	0.000051	0.000048	0.000054	0.000051	0.000053	0.000052	0.000031	0.000029	0.000046	0.000046	0.000059	0.000061	0.000030	0.000041	0.000050	0.000062	0.000032	0.000034	0.000047
Calcium (Ca)	mg/L		0.50	325	319	319	343	322	324	321	319	324	327	186	181	286	334	344	345	188	243	330	345	190	206	298
Cesium (Cs)	mg/L		0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050
Chromium (Cr)	mg/L		0.00010 to 0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050
Cobalt (Co)	mg/L		0.000050	<0.000050	<0.000050	<0.000050	<0.000050	<0.000050	<0.000050	<0.000050	<0.000050	<0.000050	<0.000050	<0.000050	<0.000050	<0.000050	<0.000050	<0.000050	<0.000050	<0.000050	<0.000050	<0.000050	<0.000050	<0.000050	<0.000050	<0.000050
Copper (Cu)	mg/L		0.000050	0.000409	0.000431	0.000405	0.000376	0.000403	0.000412	0.000387	0.000401	0.000413	0.000379	0.000518	0.000434	0.000435	0.000438	0.000405	0.000404	0.000417	0.000501	0.000423	0.000406	0.000496	0.000440	0.000456
Gallium (Ga)	mg/L		0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050
Iron (Fe)	mg/L		0.010	<0.010	<0.010	<0.010	<0.010	<0.010	<0.010	<0.010	<0.010	<0.010	<0.010	0.036	0.014	0.013	0.014	0.015	0.017	0.017	0.012	0.011	<0.010	0.023	0.014	0.014
Lead (Pb)	mg/L		0.000050	<0.000050	<0.000050	<0.000050	<0.000050	<0.000050	<0.000050	<0.000050	<0.000050	<0.000050	<0.000050	0.00186	0.000822	0.000140	0.000141	0.000388	0.000060	0.000147	0.000970	0.000122	<0.000050	0.000096	0.000052	0.000075
Lithium (Li)	mg/L		0.020	0.139	0.140	0.138	0.134	0.139	0.142	0.130	0.138	0.140	0.137	0.077	0.077	0.120	0.136	0.139</								

Appendix 3.2-1. Water Quality Data for Marine AEMP Expansion Sites, 2011

ID:				REF-Marine 2	RB1	RB1	RB1	RB1	RB1	RB2	RB2	RB2	RB2	RB2	REF-Marine 2	REF-Marine 2	REF-Marine 2	REF-Marine 2	REF-Marine 2	RB1	RB1	RB1	RB1	RB2	RB2		
Depth Zone:				Deep	Surface	Surface	Above Pycnocline	Below Pycnocline	Deep	Surface	Above Pycnocline	Above Pycnocline	Below Pycnocline	Deep	Surface	Above Pycnocline	Below Pycnocline	Below Pycnocline	Deep	Surface	Above Pycnocline	Below Pycnocline	Below Pycnocline	Deep	Surface		
Depth Sampled (m):				38	1	1	6	14	35	1	7	7	15	40	1	1	1	14	40	1	1	8	21	21	40		
Replicate:				1	1	2	1	1	1	1	1	2	1	1	1	1	1	2	1	1	1	1	2	1	2		
Date Sampled:				27-Jul-11	14-Aug-11	14-Aug-11	14-Aug-11	14-Aug-11	14-Aug-11	14-Aug-11	14-Aug-11	14-Aug-11	14-Aug-11	14-Aug-11	19-Aug-11	19-Aug-11	19-Aug-11	19-Aug-11	19-Aug-11	19-Aug-11	22-Sep-11	22-Sep-11	22-Sep-11	22-Sep-11	22-Sep-11		
ALS Sample ID:	Units	CCME Guideline for the Protection of Aquatic Life ^a	Realized Detection Limit	L1037843-4	L1044719-1	L1044719-2	L1044719-3	L1044719-7	L1044719-8	L1044719-10	L1044719-5	L1044719-6	L1044719-9	L1044719-11	L1049434-4	L1049434-3	L1049434-2	L1049434-1	L1049434-5	L1063065-1	L1063065-2	L1063065-3	L1063065-4	L1063065-5	L1063065-6	L1063065-7	
Physical Tests																											
Hardness (as CaCO ₃)	mg/L	7.0-8.7 pH g/L dependent on background levels dependent on background levels	4.3	4870	4460	4480	4790	5140	5200	4470	5010	4800	5060	5500	5000	4950	4860	5060	4920	4570	4640	4730	4810	4970	4760	4610	
pH	pH		0.10	7.83	7.93	7.92	7.91	7.88	7.82	7.94	7.90	7.92	7.89	7.82	7.88	7.85	7.86	7.86	7.81	7.87	7.85	7.83	7.82	7.79	7.87	7.86	
Salinity (EC)	g/L		1.0	26.0	23.4	23.4	25.3	27.7	28.0	24.0	26.2	25.3	27.8	28.5	26.8	27.0	27.0	27.1	27.3	26.0	26.4	27.1	27.0	27.5	26.2	26.1	
Total Suspended Solids	mg/L		2.0 to 4.0	2.5	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	
Turbidity	NTU		0.10	0.53	0.47	0.45	0.48	0.26	0.76	0.36	0.52	0.33	0.21	0.44	0.67	0.55	0.51	0.52	0.38	0.54	0.67	0.83	0.79	0.77	0.70	0.51	
Anions and Nutrients																											
Alkalinity, Total (as CaCO ₃)	mg/L	3.612 ^b	2.0	113	107	108	115	117	119	107	118	119	120	125	122	120	125	120	119	117	116	117	117	119	115	111	
Ammonia (as N)	mg/L		0.0050	0.0276	<0.0050	<0.0050	<0.0050	<0.0050	<0.0050	<0.0050	<0.0050	<0.0050	<0.0050	<0.0050	<0.0050	0.0068	0.0073	<0.0050	<0.0050	0.0161	<0.0050	<0.0050	<0.0050	<0.0050	0.0126	<0.0050	<0.0050
Bromide (Br)	mg/L		5.0	57.5	43.5	45.2	46.7	51.6	49.4	45.9	51.1	48.7	51.9	53.5	49.2	48.0	50.8	50.8	52.7	55.2	48.1	48.9	51.2	51.4	54.6	53.8	
Chloride (Cl)	mg/L		50	16200	13100	13400	14400	15100	14500	13300	14900	14300	15200	15400	14800	14100	15200	14900	15700	14800	12800	13000	14000	13500	14300	14200	
Fluoride (F)	mg/L		0.03 or 0.75	0.905	0.81	<0.75	<0.75	1.05	0.88	0.76	0.81	0.80	0.87	0.85	1.00	1.20	1.02	1.10	1.02	1.09	0.95	0.88	0.89	0.86	0.84	0.83	
Nitrate and Nitrite (as N)	mg/L		0.0060	0.0845	<0.0060	<0.0060	<0.0060	<0.0060	0.0464	<0.0060	<0.0060	<0.0060	<0.0060	<0.0060	0.0685	<0.0060	<0.0060	0.0174	0.0214	<0.0060	<0.0060	0.0113	<0.0060	<0.0060	0.0458	<0.0060	<0.0060
Nitrate (as N)	mg/L		0.0060	0.0845	<0.0060	<0.0060	<0.0060	<0.0060	0.0464	<0.0060	<0.0060	<0.0060	<0.0060	<0.0060	0.0685	<0.0060	<0.0060	0.0152	0.0191	<0.0060	<0.0060	0.0113	<0.0060	<0.0060	0.0458	<0.0060	<0.0060
Nitrite (as N)	mg/L		0.0020	<0.0020	<0.0020	<0.0020	<0.0020	<0.0020	<0.0020	<0.0020	<0.0020	<0.0020	<0.0020	<0.0020	<0.0020	<0.0020	<0.0020	0.0022	0.0023	<0.0020	<0.0020	<0.0020	<0.0020	<0.0020	<0.0020	<0.0020	
Total Phosphorus	mg/L		0.0020	0.0566	0.0285	0.0308	0.0336	0.0342	0.0564	0.0294	0.0347	0.0320	0.0344	0.0484	0.0442	0.0462	0.0467	0.0457	0.0685	0.0321	0.0344	0.0378	0.0387	0.0461	0.0323	0.0322	
Sulphate (SO ₄)	mg/L		50	2170	1860	1900	2030	2140	2060	1900	2110	2020	2160	2180	2110	1990	2160	2120	2230	2110	1820	1840	1990	1910	2020	2020	
Cyanides																											
Total Cyanide	mg/L		0.0010	0.0010	<0.0010	<0.0010	<0.0010	<0.0010	<0.0010	<0.0010	<0.0010	0.0012	<0.0010	0.0013	0.0015	0.0015	0.0015	0.0016	0.0015	0.0013	0.0013	0.0012	0.0010	<0.0010	0.0016	0.0014	
Free Cyanide	mg/L		0.0010 or 0.0050	<0.0050	<0.0010	<0.0010	<0.0010	<0.0010	<0.0010	0.0011	<0.0010	<0.0010	0.0011	0.0010	<0.0010	<0.0010	<0.0010	<0.0010	<0.0010	<0.0010	<0.0010	<0.0010	<0.0010	<0.0010	<0.0010	<0.0010	
Organic / Inorganic Carbon																											
Total Organic Carbon	mg/L		0.50	1.27	1.83	1.73	1.59	1.49	1.60	1.72	1.51	1.54	1.52	1.53	1.51	1.43	1.43	1.59	1.31	1.71	1.71	1.60	1.60	1.54	1.68	1.61	
Total Metals																											
Aluminum (Al)	mg/L	0.0125 ^b	0.0050	0.0127	0.0064	0.0085	<0.0050	0.0096	0.0053	<0.0050	<0.0050	<0.0050	<0.0050	<0.0050	<0.0050	<0.0050	0.0056	0.0188	<0.0050	0.0072	0.0091	0.0225	0.0254	0.0168	0.0062	0.0122	
Antimony (Sb)	mg/L		0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	
Arsenic (As)	mg/L		0.0004	0.00125	0.00095	0.00106	0.00112	0.00090	0.00115	0.00053	0.00111	0.00098	0.00115	0.00124	0.00107	0.00095	0.00108	0.00123	0.00097	0.00088	0.00093	0.00083	0.00098	0.00094	0.00087	0.00083	
Barium (Ba)	mg/L		0.0010	0.0129	0.0101	0.0099	0.0111	0.0126	0.0122	0.0106	0.0119	0.0113	0.0118	0.0121	0.0121	0.0119	0.0122	0.0118	0.0133	0.0061	0.0069	0.0139	0.0130	0.0136	0.0063	0.0069	
Beryllium (Be)	mg/L		0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	
Bismuth (Bi)	mg/L		0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	
Boron (B)	mg/L		0.10	3.77	3.22	3.22	3.56	3.79	3.82	3.23	3.66	3.53	3.68	3.73	3.81	3.81	3.90	3.76	3.86	3.22	3.40	3.55	3.48	3.61	3.33	3.49	
Cadmium (Cd)	mg/L		0.000020	0.000062	0.000039	0.000036	0.000041	0.000044	0.000055	0.000065	0.000045	0.000041	0.000039	0.000056	0.000048	0.000051	0.000052	0.000053	0.000076	0.000049</							

Appendix 3.2-1. Water Quality Data for Marine AEMP Expansion Sites, 2011

Site ID:				RB2	RB2	RB2	REF-Marine 2	REF-Marine 2	REF-Marine 2	REF-Marine 2
Depth Zone:				Above	Below	Deep	Surface	Above	Below	Deep
Depth Sampled (m):				Pycnocline	Pycnocline			Pycnocline	Pycnocline	
Replicate:				1	21	40	1	8	20	40
Date Sampled:				1	1	1	1	1	1	1
ALS Sample ID:	Units	CCME Guideline for the Protection of Aquatic Life ^a	Realized Detection Limit	22-Sep-11 L1063065-8	22-Sep-11 L1063065-9	22-Sep-11 L1063065-10	25-Sep-11 L1064757-1	25-Sep-11 L1064757-2	25-Sep-11 L1064757-3	25-Sep-11 L1064757-4
Physical Tests										
Hardness (as CaCO ₃)	mg/L	7.0-8.7 <10% fluctuation ^b dependent on background levels dependent on background levels	4.3	4750	4770	4850	4520	5030	4910	5080
pH	pH		0.10	7.86	7.84	7.80	7.99	8.00	8.00	7.98
Salinity (EC)	g/L		1.0	26.5	26.8	27.4	23.6	26.0	26.3	27.4
Total Suspended Solids	mg/L		2.0 to 4.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0
Turbidity	NTU		0.10	0.55	0.69	0.40	1.14	0.52	0.60	0.47
Anions and Nutrients										
Alkalinity, Total (as CaCO ₃)	mg/L	3.612 ^b	2.0	115	115	119	105	110	110	116
Ammonia (as N)	mg/L		0.0050	0.0055	<0.0050	0.0144	<0.0050	<0.0050	<0.0050	0.0499
Bromide (Br)	mg/L		5.0	55.2	52.8	53.1	47.7	52.8	51.7	52.9
Chloride (Cl)	mg/L		50	14700	14000	14900	13800	14800	14800	15100
Fluoride (F)	mg/L		0.03 or 0.75	0.94	0.83	0.86	1.04	0.98	0.92	0.82
Nitrate and Nitrite (as N)	mg/L		0.0060	<0.0060	<0.0060	0.0629	<0.0060	<0.0060	<0.0060	0.0669
Nitrate (as N)	mg/L		0.0060	<0.0060	<0.0060	0.0629	<0.0060	<0.0060	<0.0060	0.0628
Nitrite (as N)	mg/L		0.0020	<0.0020	<0.0020	<0.0020	<0.0020	<0.0020	<0.0020	0.0041
Total Phosphorus	mg/L		0.0020	0.0342	0.0366	0.0468	0.0354	0.0342	0.0336	0.0491
Sulphate (SO ₄)	mg/L		50	2090	2000	2120	1970	2110	2120	2160
Cyanides										
Total Cyanide	mg/L		0.0010	0.0014	0.0013	0.0012	<0.0010	<0.0010	<0.0010	<0.0010
Free Cyanide	mg/L		0.0010 or 0.0050	<0.0010	<0.0010	<0.0010	<0.0010	<0.0010	<0.0010	<0.0010
Organic / Inorganic Carbon										
Total Organic Carbon	mg/L		0.50	1.69	1.57	1.46	1.88	1.38	1.37	1.15
Total Metals										
Aluminum (Al)	mg/L	0.0125 ^b	0.0050	0.0087	0.0121	0.0092	0.0355	0.0122	0.0078	0.0057
Antimony (Sb)	mg/L		0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050
Arsenic (As)	mg/L		0.0004	0.00086	0.00089	0.00095	0.00100	0.00115	0.00101	0.00108
Barium (Ba)	mg/L		0.0010	0.0076	0.0115	0.0115	0.0039	0.0077	0.0077	0.0119
Beryllium (Be)	mg/L		0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050
Bismuth (Bi)	mg/L	0.00012	0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050
Boron (B)	mg/L		0.10	3.51	3.70	3.70	3.18	3.61	3.60	3.70
Cadmium (Cd)	mg/L		0.000020	0.000055	0.000052	0.000061	0.000033	0.000038	0.000035	0.000048
Calcium (Ca)	mg/L		0.50	323	323	327	301	332	323	328
Cesium (Cs)	mg/L		0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050
Chromium (Cr)	mg/L		0.00010 to 0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050
Cobalt (Co)	mg/L		0.000050	<0.000050	<0.000050	<0.000050	<0.000050	<0.000050	<0.000050	<0.000050
Copper (Cu)	mg/L		0.000050	0.000459	0.000458	0.000389	0.000431	0.000359	0.000340	0.000327
Gallium (Ga)	mg/L		0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050
Iron (Fe)	mg/L		0.010	0.014	0.019	0.014	0.041	0.012	<0.010	0.011
Lead (Pb)	mg/L	Inorganic Hg: 0.000016 ^b Inorganic Hg: 0.016 ^b	0.000050	0.000090	0.000137	<0.000050	<0.000050	<0.000050	<0.000050	<0.000050
Lithium (Li)	mg/L		0.020	0.130	0.136	0.136	0.123	0.140	0.139	0.142
Magnesium (Mg)	mg/L		1.0	957	962	981	916	1020	997	1030
Manganese (Mn)	mg/L		0.000050	0.00170	0.00164	0.00177	0.00182	0.00128	0.00118	0.00277
Mercury (Hg)	mg/L		0.000010	-	-	-	-	-	-	-
Mercury (Hg) (ultra low-level)	µg/L		0.00050	0.00118	0.00177	0.00126	0.00070	0.00077	0.00069	0.00068
Molybdenum (Mo)	mg/L		0.0020	0.0087	0.0093	0.0091	0.0072	0.0081	0.0080	0.0082
Nickel (Ni)	mg/L		0.000050	0.000569	0.000581	0.000578	0.000444	0.000452	0.000443	0.000483
Phosphorus (P)	mg/L		1.0	<1.0	<1.0	<1.0	<1.0	<1.0	<1.0	<1.0
Potassium (K)	mg/L		20	300	298	304	285	315	310	318
Rhenium (Re)	mg/L	0.00010 or 0.00020	0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050
Rubidium (Rb)	mg/L		0.0050	0.0873	0.0920	0.0948	0.0796	0.0893	0.0876	0.0902
Selenium (Se)	mg/L		0.00040	<0.00040	<0.00040	<0.00040	<0.00040	<0.00040	<0.00040	<0.00040
Silicon (Si)	mg/L		0.50	<0.50	<0.50	0.66	<0.50	<0.50	<0.50	0.72
Silver (Ag)	mg/L		0.00010 or 0.00020	<0.00010	<0.00010	<0.00010	<0.00010	<0.00010	<0.00010	<0.00010
Sodium (Na)	mg/L		20	8570	8570	8690	8120	8950	8790	9010
Strontium (Sr)	mg/L		0.010 or 0.050	5.83	5.78	5.89	5.59	6.17	6.04	6.16
Tellurium (Te)	mg/L		0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050
Thallium (Tl)	mg/L		0.000050	<0.000050	<0.000050	<0.000050	<0.000050	<0.000050	<0.000050	<0.000050
Thorium (Th)	mg/L		0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050
Tin (Sn)	mg/L	0.0010 or 0.0020	0.0010 or 0.0020	<0.0010	<0.0010	<0.0010	<0.0020	<0.0020	<0.0020	<0.0020
Titanium (Ti)	mg/L		0.0050	<0.0050	<0.0050	<0.0050	<0.0050	<0.0050	<0.0050	<0.0050
Tungsten (W)	mg/L		0.0010	<0.0010	<0.0010	<0.0010	<0.0010	<0.0010	<0.0010	<0.0010
Uranium (U)	mg/L		0.000050	0.00217	0.00212	0.00230	0.00208	0.00227	0.00234	0.00233
Vanadium (V)	mg/L		0.00050	0.00079	0.00077	0.00084	0.00078	0.00077	0.00073	0.00076
Yttrium (Y)	mg/L		0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050
Zinc (Zn)	mg/L		0.00080	<0.00080	<0.00080	<0.00080	<0.00080	<0.00080	<0.00080	<0.00080
Zirconium (Zr)	mg/L		0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050
Speciated Metals										
Methylmercury (Total)	µg/L		0.000050	<0.000050	<0.000050	<0.000050	<0.000050	<0.000050	<0.000050	<0.000050
Radiochemistry										
Radium-226	Bq/L		0.005	<0.005	<0.005	<0.005	<0.005	<0.005	<0.005	0.01

Notes:

a) Canadian water quality guidelines for the protection of marine aquatic life, Canadian Council of Ministers of the Environment, Updated January 2011

b) Interim guideline

Appendix 3.2-2

**QA/QC Blank Data for Marine AEMP Expansion Water
Quality Sampling, Doris North Project, 2011**

Appendix 3.2-2. QA/QC Blank Data for Marine AEMP Expansion Water Quality Sampling, 2011

Blank Type: Date Sampled: ALS Sample ID:	Units	Realized Detection Limit	Field Blank 24-Apr-11 L998223-14	Equipment Blank 21-Jul-11 L1035341-5	Field Blank 22-Jul-11 L1035865-15	Equipment Blank 14-Aug-11 L1044719-4	Equipment Blank 20-Sep-11 L1062230-5	Field Blank 20-Sep-11 L1062230-6	Travel Blank 20-Sep-11 L1062230-7
Physical Tests									
Hardness (as CaCO ₃)	mg/L	0.50	<0.50	<0.50	<0.50	<0.50	1.97	<0.50	<0.50
pH	pH	0.10	5.63	4.37	5.93	5.51	3.14	5.39	5.60
Salinity (EC)	g/L	1.0	<1.0	<1.0	<1.0	<1.0	<1.0	<1.0	<1.0
Total Suspended Solids	mg/L	2.0	<2.0	2.0	<2.0	<2.0	<2.0	<2.0	<2.0
Turbidity	NTU	0.10	<0.10	0.71	<0.10	<0.10	0.10	<0.10	<0.10
Anions and Nutrients									
Alkalinity, Total (as CaCO ₃)	mg/L	2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0
Ammonia (as N)	mg/L	0.0050	<0.0050	0.0376	<0.0050	<0.0050	0.0231	<0.0050	<0.0050
Bromide (Br)	mg/L	0.050 or 0.10	<0.050	<0.050	<0.10	<0.050	<0.050	<0.050	<0.050
Chloride (Cl)	mg/L	0.50	<0.50	2.75	<0.50	<0.50	2.44	<0.50	<0.50
Fluoride (F)	mg/L	0.20	<0.020	<0.020	<0.020	<0.020	<0.020	<0.020	<0.020
Nitrate and Nitrite (as N)	mg/L	0.0060	<0.0060	<0.0060	<0.0060	0.121	13.1	<0.0060	<0.0060
Nitrate (as N)	mg/L	0.0050 or 0.0060	<0.0060	<0.0050	<0.0060	0.121	13.1	<0.0060	<0.0060
Nitrite (as N)	mg/L	0.0020	<0.0020	<0.0010	<0.0020	<0.0020	0.0079	<0.0020	<0.0020
Total Phosphorus	mg/L	0.0020	<0.0020	<0.0020	<0.0020	<0.0020	<0.0020	<0.0020	<0.0020
Sulphate (SO ₄)	mg/L	0.50	<0.50	<0.50	<0.50	<0.50	<0.50	<0.50	<0.50
Cyanides									
Total Cyanide	mg/L	0.0010 to 0.00125	<0.0010	0.0127	<0.0010	<0.0010	0.204	<0.0011	<0.0010
Free Cyanide	mg/L	0.0010 or 0.0050	<0.0050	<0.0050	<0.0050	<0.0010	<0.0010	<0.0010	<0.0010
Organic / Inorganic Carbon									
Total Organic Carbon	mg/L	0.50	<0.50	0.99	<0.50	0.64	7.85	<0.50	<0.50
Total Metals									
Aluminum (Al)	mg/L	0.0050	<0.0050	0.0467	<0.0050	<0.0050	0.153	<0.0050	<0.0050
Antimony (Sb)	mg/L	0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050
Arsenic (As)	mg/L	0.00040	<0.00040	<0.00040	<0.00040	<0.00040	<0.00040	<0.00040	<0.00040
Barium (Ba)	mg/L	0.0010	<0.0010	<0.0010	<0.0010	0.0084	0.0479	<0.0010	<0.0010
Beryllium (Be)	mg/L	0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050
Bismuth (Bi)	mg/L	0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050
Boron (B)	mg/L	0.10	<0.10	<0.10	<0.10	<0.10	<0.10	<0.10	<0.10
Cadmium (Cd)	mg/L	0.000020	<0.000020	<0.000020	<0.000020	<0.000020	<0.000020	<0.000020	<0.000020
Calcium (Ca)	mg/L	0.050	<0.050	<0.050	<0.050	<0.050	0.478	<0.050	<0.050
Cesium (Cs)	mg/L	0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050
Chromium (Cr)	mg/L	0.00010 or 0.00050	<0.00010	0.0459	<0.00050	<0.00010	<0.00050	<0.00050	<0.00050
Cobalt (Co)	mg/L	0.000050	<0.000050	0.00166	<0.000050	<0.000050	0.000051	<0.000050	<0.000050
Copper (Cu)	mg/L	0.000050	<0.000050	0.00128	<0.000050	0.000080	0.00314	<0.000050	<0.000050
Gallium (Ga)	mg/L	0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050
Iron (Fe)	mg/L	0.010	<0.010	0.130	<0.010	<0.010	0.020	<0.010	<0.010
Lead (Pb)	mg/L	0.000050	<0.000050	0.000061	<0.000050	<0.000050	0.000625	<0.000050	<0.000050
Lithium (Li)	mg/L	0.020	<0.020	<0.020	<0.020	<0.020	<0.020	<0.020	<0.020
Magnesium (Mg)	mg/L	0.10	<0.10	<0.10	<0.10	<0.10	0.19	<0.10	<0.10
Manganese (Mn)	mg/L	0.000050	<0.000050	0.00360	<0.000050	0.000212	0.000581	<0.000050	<0.000050
Mercury (Hg)	mg/L	0.000010	<0.000010	-	-	-	-	-	-
Mercury (Hg) (ultra low-level)	µg/L	0.00050	-	0.00060	<0.00050	<0.00050	0.00211	<0.00050	<0.00050
Molybdenum (Mo)	mg/L	0.0020	<0.0020	<0.0020	<0.0020	<0.0020	<0.0020	<0.0020	<0.0020
Nickel (Ni)	mg/L	0.000050	<0.000050	0.0486	<0.000050	<0.000050	0.000298	<0.000050	<0.000050
Phosphorus (P)	mg/L	1.00	<1.0	<1.0	<1.0	<1.0	<1.0	<1.0	<1.0
Potassium (K)	mg/L	2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0
Rhenium (Re)	mg/L	0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050
Rubidium (Rb)	mg/L	0.0050	<0.0050	<0.0050	<0.0050	<0.0050	<0.0050	<0.0050	<0.0050
Selenium (Se)	mg/L	0.00040	<0.00040	<0.00040	<0.00040	<0.00040	<0.00040	<0.00040	<0.00040
Silicon (Si)	mg/L	0.050	<0.050	0.085	<0.050	<0.050	<0.050	<0.050	<0.050
Silver (Ag)	mg/L	0.00010	<0.00010	<0.00010	<0.00010	<0.00010	<0.00010	<0.00010	<0.00010
Sodium (Na)	mg/L	2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0
Strontium (Sr)	mg/L	0.0050 or 0.010	<0.010	<0.0050	<0.0050	<0.0050	<0.0050	<0.0050	<0.0050
Tellurium (Te)	mg/L	0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050
Thallium (Tl)	mg/L	0.000050	<0.000050	<0.000050	<0.000050	<0.000050	<0.000050	<0.000050	<0.000050
Thorium (Th)	mg/L	0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050
Tin (Sn)	mg/L	0.0010	<0.0010	<0.0010	<0.0010	<0.0010	0.0089	<0.0010	<0.0010
Titanium (Ti)	mg/L	0.0050	<0.0050	<0.0050	<0.0050	<0.0050	<0.0050	<0.0050	<0.0050
Tungsten (W)	mg/L	0.0010	<0.0010	<0.0010	<0.0010	<0.0010	<0.0010	<0.0010	<0.0010
Uranium (U)	mg/L	0.000050	<0.000050	<0.000050	<0.000050	<0.000050	<0.000050	<0.000050	<0.000050
Vanadium (V)	mg/L	0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050
Yttrium (Y)	mg/L	0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050
Zinc (Zn)	mg/L	0.00080	<0.00080	0.0329	<0.00080	<0.00080	0.00985	<0.00080	<0.00080
Zirconium (Zr)	mg/L	0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050	<0.00050
Speciated Metals									
Methylmercury (Total)	µg/L	0.000050	-	<0.000050	<0.000050	<0.000050	0.000328	<0.000050	<0.000050
Radiochemistry									
Radium-226	Bq/L	0.005	-	<0.005	-	<0.006	<0.005	-	-

Note:

Bold values represent concentrations that are higher than analytical detection limits.

Appendix 3.3-1

**Sediment Quality Data for Marine AEMP Expansion Sites,
Doris North Project, 2011**

Appendix 3.3-1. Sediment Quality Data for Marine AEMP Expansion Sites, Doris North Project, 2011

Site ID: Depth Sampled (m): Replicate: Date Sampled: ALS Sample ID:		CCME Guidelines for the Protection of Aquatic Life ^a		Realized Detection Limit	RB1 42 1 23-Aug-11 L1050746-5	RB1 42 2 23-Aug-11 L1050746-6	RB1 42 3 23-Aug-11 L1050746-7	REF-Marine 2 42 1 24-Aug-11 L1050746-2	REF-Marine 2 42 2 24-Aug-11 L1050746-3	REF-Marine 2 42 3 24-Aug-11 L1050746-4
	Units	ISQG ^b	PEL ^c							
Physical Tests										
Moisture	%			0.25	57.6	55.6	58.3	53.7	57.8	57.9
pH	pH			0.10	7.29	7.28	7.21	7.48	7.28	7.31
Particle Size										
% Gravel (>2 mm)	%			0.10	<0.10	<0.10	12.4	<0.10	<0.10	<0.10
% Sand (2.0 mm - 0.063 mm)	%			0.10	2.63	2.08	3.05	2.16	1.85	3.16
% Silt (0.063 mm - 4 µm)	%			0.10	57.6	47.1	46.4	56.8	60.0	57.0
% Clay (<4 µm)	%			0.10	39.8	50.9	38.1	41.0	38.2	39.8
Anions & Nutrients										
Total Nitrogen	%			0.020	0.111	0.112	0.127	0.133	0.130	0.135
Organic / Inorganic Carbon										
Total Organic Carbon	%			0.10	0.80	0.72	0.74	1.07	1.05	1.21
Plant Available Nutrients										
Available Ammonium-N	mg/kg			1.6 or 2.4	16.9	29.8	24.7	17.9	11.4	12.4
Available Nitrate-N	mg/kg			4.0 or 6.0	<6.0	<4.0	<6.0	<4.0	<6.0	<4.0
Available Nitrite-N	mg/kg			0.80 or 1.2	<1.2	<0.80	<1.2	<0.80	<1.2	<0.80
Available Phosphate-P	mg/kg			4.0 or 6.0	73.3	91.5	101	88.6	81.5	103
Metals										
Aluminum (Al)	mg/kg			50	20700	22100	21200	15600	16200	15700
Antimony (Sb)	mg/kg			0.10	0.24	0.23	0.24	0.29	0.30	0.31
Arsenic (As)	mg/kg	7.24	41.6	0.050	51.9	13.6	45.0	51.0	47.1	55.8
Barium (Ba)	mg/kg			0.50	148	142	147	107	111	113
Beryllium (Be)	mg/kg			0.20	0.64	0.65	0.64	0.53	0.53	0.50
Bismuth (Bi)	mg/kg			0.20	<0.20	0.28	<0.20	<0.20	<0.20	<0.20
Cadmium (Cd)	mg/kg	0.7	4.2	0.050	0.126	0.135	0.107	0.095	0.104	0.086
Calcium (Ca)	mg/kg			50	7250	6820	7280	7710	6910	7290
Chromium (Cr)	mg/kg	52.3	160	0.50	64.1	66.3	64.3	46.3	48.6	46.9
Cobalt (Co)	mg/kg			0.10	13.1	12.9	13.0	8.60	9.06	8.85
Copper (Cu)	mg/kg	18.7	108	0.50	24.2	25.9	23.8	18.4	19.7	19.1
Iron (Fe)	mg/kg			50	49900	36400	48800	36600	39200	42000
Lead (Pb)	mg/kg	30.2	112	0.50	8.23	8.24	8.32	5.98	6.34	6.28
Lithium (Li)	mg/kg			1.0	41.9	46.7	44.9	31.1	32.1	30.4
Magnesium (Mg)	mg/kg			20	17100	17800	17500	12200	13300	12800
Manganese (Mn)	mg/kg			1.0	477	384	444	270	299	302
Mercury (Hg)	mg/kg	0.13	0.70	0.0050 or 0.015	0.0152	0.0145	0.0148	0.016	0.019	0.020
Molybdenum (Mo)	mg/kg			0.50	1.73	1.64	1.61	1.33	1.27	1.23
Nickel (Ni)	mg/kg			0.50	31.5	32.6	31.3	21.8	23.1	22.2
Phosphorus (P)	mg/kg			50	4640	1610	4200	5050	4850	6000
Potassium (K)	mg/kg			100	7750	7790	7790	5960	6230	6140
Selenium (Se)	mg/kg			0.20	0.57	0.49	0.51	0.59	0.62	0.63
Silver (Ag)	mg/kg			0.10	<0.10	<0.10	<0.10	<0.10	0.10	<0.10
Sodium (Na)	mg/kg			100	12600	12800	13400	10100	13700	12100
Strontium (Sr)	mg/kg			0.50	139	67.3	122	169	150	180
Sulphur (S)	mg/kg			500	1500	1700	1600	1400	1400	1600
Thallium (Tl)	mg/kg			0.050	0.265	0.270	0.259	0.183	0.199	0.192
Tin (Sn)	mg/kg			2.0	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0
Titanium (Ti)	mg/kg			1.0	1330	1540	1390	1020	1060	1010
Uranium (U)	mg/kg			0.050	2.28	2.25	2.14	2.18	2.13	2.01
Vanadium (V)	mg/kg			0.20	85.5	82.1	84.6	62.9	67.2	66.2
Zinc (Zn)	mg/kg	124	271	1.0	75.1	78.1	74.8	54.0	56.1	54.1
Speciated Metals										
Methylmercury	mg/kg			0.000050	<0.000050	<0.000050	<0.000050	<0.000050	0.000061	<0.000050

Notes:

Shaded cells indicate values that exceed CCME guidelines.

a) Canadian sediment quality guidelines for the protection of marine aquatic life, Canadian Council of Ministers of the Environment, Updated January 2011

b) ISQG = Interim Sediment Quality Guideline

c) PEL = Probable Effects Level

Appendix 3.4-1

**Phytoplankton Biomass (as Chlorophyll a) Data for Marine
AEMP Expansion Sites, Doris North Project, 2011**

Appendix 3.4-1. Phytoplankton Biomass (as Chlorophyll *a*) Data for Marine AEMP Expansion Sites, Doris North Project, 2011

Marine Site	Date Sampled	Depth Sampled	ALS Sample ID	Replicate #	Phytoplankton Biomass		
					(µg chl <i>a</i> /L)	Mean	SE
RB1	23-Apr-11	3 m	L1012910-22	1	0.130	0.136	0.008
RB1	23-Apr-11	3 m	L1012910-23	2	0.152		
RB1	23-Apr-11	3 m	L1012910-24	3	0.125		
RB1	23-Jul-11	1 m	L1047581-36	1	0.164	0.144	0.018
RB1	23-Jul-11	1 m	L1047581-37	2	0.160		
RB1	23-Jul-11	1 m	L1047581-38	3	0.107		
RB1	14-Aug-11	1 m	L1052615-7	1	0.173	0.176	0.024
RB1	14-Aug-11	1 m	L1052615-8	2	0.220		
RB1	14-Aug-11	1 m	L1052615-9	3	0.136		
RB1	22-Sep-11	1 m	L1066523-1	1	0.270	0.422	0.156
RB1	22-Sep-11	1 m	L1066523-2	2	0.733		
RB1	22-Sep-11	1 m	L1066523-3	3	0.263		
RB2	23-Apr-11	3 m	L1012910-19	1	0.231	0.179	0.030
RB2	23-Apr-11	3 m	L1012910-20	2	0.178		
RB2	23-Apr-11	3 m	L1012910-21	3	0.128		
RB2	23-Jul-11	1 m	L1047581-39	1	0.120	0.091	0.038
RB2	23-Jul-11	1 m	L1047581-40	2	0.138		
RB2	23-Jul-11	1 m	L1047581-41	3	0.016		
RB2	14-Aug-11	1 m	L1052615-10	1	0.219	0.228	0.019
RB2	14-Aug-11	1 m	L1052615-11	2	0.264		
RB2	14-Aug-11	1 m	L1052615-12	3	0.200		
RB2	22-Sep-11	1 m	L1066523-4	1	0.663	0.499	0.125
RB2	22-Sep-11	1 m	L1066523-5	2	0.582		
RB2	22-Sep-11	1 m	L1066523-6	3	0.253		
REF-Marine 2	21-Jul-11	1 m	L1047581-45	1	0.167	0.189	0.011
REF-Marine 2	21-Jul-11	1 m	L1047581-46	2	0.204		
REF-Marine 2	21-Jul-11	1 m	L1047581-47	3	0.196		
REF-Marine 2	19-Aug-11	1 m	L1052615-25	1	1.18	1.50	0.32
REF-Marine 2	19-Aug-11	1 m	L1052615-26	2	1.19		
REF-Marine 2	19-Aug-11	1 m	L1052615-27	3	2.14		
REF-Marine 2	25-Sep-11	1 m	L1066523-7	1	0.570	0.627	0.080
REF-Marine 2	25-Sep-11	1 m	L1066523-8	2	0.527		
REF-Marine 2	25-Sep-11	1 m	L1066523-9	3	0.785		

SE = standard error of the mean

Appendix 3.5-1

**Benthos Taxonomy Data for Marine AEMP Expansion Sites,
Doris North Project, 2011**

Appendix 3.5-1. Benthos Taxonomy Data for Marine AEMP Expansion Sites, Doris North Project, 2011

Taxonomic Identification			RB1 23-Aug-11					REF-Marine 2 24-Aug-11										
Major Group	Family	Genus/Species	Rep-1		Rep-2		Rep-3		Rep-4		Rep-5							
			A	J	A	J	A	J	A	J	A	J	A	J				
CNIDARIA																		
Anthozoa	Actiniaria (Order; symbiont on Colus sp.)*	Actiniaria Indet.*							1			1						
	Actiniaria (Order)*	Actiniaria Indet.*								1								
NEMERTEA																		
	Lineidae	Cerebratulus sp.		1														
		Lineidae Indet.					1		2	1	2	1	3					
	Hoplonemertea (Order)*	Hoplonemertea Indet.*			1		1											
	Tubulanidae	Tubulanus sp.											1					
ANNELIDA																		
Polychaeta Errantia	Polynoidae	Melaenis loveni										1						
		Polynoidae Indet.							2			2						
	Hesionidae	Hesionidae Indet.			2	4	7	1	4	1	4	1	2					
	Lumbrineridae	Lumbrineridae Indet.			1		1											
		Lumbrineris fragilis											1					
		Lumbrineris sp.							3	1	3	3	5	1				
	Nephtyidae	Nephtys ciliata		3	2		1						1					
		Nephtys ferruginea							1	2		1						
		Nephtys sp.	1	1	3	1		4	5	19	2	12						
	Hesionidae	Nereimyra sp.				2				3	1	5						
	Pholoidae	Pholoe nr. minuta							1	2	2	1						
		Pholoe sp.		1	1	1	3	2		1	1		3					
Polychaeta Sedentaria	Ampharetidae	Ampharete sp.							2									
	Cirratulidae	Aphelochoeta sp.		4	4	7	12	1	7	24	1	26	13	22	1			
		Chaetozone sp.						2	1			8	2	7				
		Cirratulidae Indet.				6	4		11			2	18	11	18			
	Paraonidae	Aricidea sp.					11			1		6		6				
		Paraonidae Indet.		1								1						
		Levinsonia gracilis			1		1											
	Capitellidae	Capitella capitata Cmplx.							1				1	1				
		Capitellidae Indet.									1			1				
		Mediomastus sp.		1			3	3	3	3	3	9	3	10				
	Sabellidae	Chone sp.						1	1				1					
	Cossuridae	Cossura sp.												1				
	Maldanidae	Euclymeninae Indet.						1										
		Maldane sarsi	20	8	16	4	17	3	11	32	1	14	1	19	1	11	2	18
	Orbiniidae	Leitoscoloplos sp.		3	2	2	2	2	2	4	2	2	1	3				
	Spionidae	Prionospio nr. cirrifera							1									
		Prionospio sp.								3	5	3	3	2	7			
	Ampharetidae	Melinna sp.						1										
	Pectinariidae	Pectinaria granulata							1									
	Scalibregmatidae	Scalibregma inflatum										1		1	2			
	Chaetopteridae	Spiochaetopterus typicus		1														
	Terebellidae	Streblosoma sp.													1			
		Terebellidae Indet.						1										
Trichobranchidae	Terebellides sp.		3	1	1	3	2		4		1							
ARTHROPODA																		
Amphipoda	Lysianassidae	Anonyx sp.	1															
		Boeckosimus affinis	7	4	7	1	2	4	2	1	2	1	1	1				
	Corophiidae	Corophium sp.							1									
	Dexaminidae	Guernea nordenskioldi			1													
	Ampeliscidae	Haploops nr. laevis		1	4	1	2											
		Haploops sp.								1	2		1					
Pontoporeiidae	Pontoporeia femorata							1										
Ostracoda	Ostracoda (Class)	Ostracoda Indet.	2		8	6	20			1	1	2						

Appendix 3.5-1. Benthos Taxonomy Data for Marine AEMP Expansion Sites, Doris North Project, 2011

Taxonomic Identification			RB1 23-Aug-11					REF-Marine 2 24-Aug-11				
Major Group	Family	Genus/Species	Rep-1		Rep-2		Rep-3		Rep-4		Rep-5	
			A	J	A	J	A	J	A	J	A	J
Cumacea	Diastylidae	<i>Brachydiastylis resima</i>	3		5		14		14		33	
		<i>Diastylis</i> nr. <i>dalli</i>										
		<i>Diastylis</i> nr. <i>goodsiri</i>										
		<i>Diastylis rathkei</i>							1		1	
		<i>Diastylis</i> sp. (with spines)									1	
		<i>Diastylis</i> sp. (no spines)			1							
	Leuconidae	<i>Eudorella</i> sp.						1	1		2	
		<i>Eudorellopsis</i> sp.						2		1	2	
		<i>Leucon</i> sp.										1
Isopoda	Gnathiidae	<i>Gnathia</i> sp.				1		1				1
Tanaidacea	Leptocheliidae	<i>Leptochelia</i> sp.								2		
	Tanaidacea (Order)*	Tanaidacea Indet.*						1				
MOLLUSCA												
Aplacophora	Aplacophora (Class)*	Aplacophora Indet.*						1				
Gastropoda	Cylichnidae	<i>Cylichna alba</i>	2							1	1	
	Buccinidae	<i>Colus</i> sp. (with anemone symbiont)							1			1
	Haminoeidae	<i>Haminoea</i> sp.										1
Bivalvia	Astartidae	<i>Astarte borealis</i>	8	1		7		7	2			
	Thyasiridae	<i>Axinopsida orbiculata</i>							1			
	Tellinidae	<i>Macoma calcareea</i>										1
	Yoldiidae	<i>Portlandia arctica</i>				1						
ECHINODERMATA												
Asteroidea	Asteriidae	<i>Leptasterias</i> sp.								1		
Ophiuroidea	Ophiuridae	<i>Ophiura</i> sp.	1		3		1	3	4			
UROCHORDATA												
Ascidiacea	Corellidae	<i>Chelyosoma</i> sp.			1							
TOTAL number of individuals			45	1	42	0	77	11	68	7	134	10
									84	9	84	5
											140	2
											67	7
											150	4

Notes:

* Taxa marked with an asterisk were excluded from total counts and from all benthos analyses because these taxa were not identifiable to the family level and made up a minor proportion (<2%) of the benthos assemblage in each replicate.

The total number of individuals was divided by 3 times the surface area of the Petite Ponar (i.e., 3 x 0.023 m ²) to determine the benthos density in units of organisms/m ² (because each replicate consisted of 3 pooled subsample Ponar grabs).

A = adult

J = juvenile