

BIBLIOGRAPHY

- Armitage, A.E., Tella, S. and Miller, A.R. (1993) Iron-formation hosted gold mineralization and its geological setting, Meliadine Lake area, District of Keewatin, Northwest Territories; in Current Research, Geological Survey of Canada, 1993-1C, p. 187-195.
- Besserer, D.J., 1994. Regional Metamorphism and P-T conditions of the Gibson-MacQuoid Lake Greenstone Belt, District of Keewatin, N.W.T. Unpublished undergraduate paper submitted in partial fulfilment of the requirements for Geology 490, The University of Western Ontario.
- Comaplex Minerals Corporation (Comaplex), 2002. Press Release titled 'Meliadine West Project; \$500,000 Option Payment Received'. January 18, 2002 Public Press Release.
- Cumberland Resources Ltd.(Cumberland), 2002. Press Release titled 'Meadowbank Gold Project: Revised Economic Studies Indicate Improved Economics. Major Program Planned For 2002' January 24, 2002 Public Press Release.
- Dawson, J.B., and Stephens, W.E. (1975) Statistical classification of garnets from kimberlite and associated xenoliths. *Journal of Geology*, Vol.83, p.589-607.
- Dickson, G. (1991) Geology and gold mineralization on the Asamera/Comaplex Meliadine Project, Rankin Inlet area, N.W.T. (abstract); in *Exploration Overview 1990, Northwest Territories*; (ed.) S.P. Goff.
- Fipke, C.E., Gurney, J.J. and Moore, R.O. (1995) Diamond exploration techniques emphasising indicator mineral geochemistry and Canadian examples; *Geological Survey of Canada, Bulletin 423*, 86 pages.
- Gurney, J.J., (1984) A correlation between garnets and diamonds in kimberlite. *In Kimberlite Occurrence and Origin: A basis for conceptual models in exploration*, J.E. Glover and P.G. Harris (eds.). Geology Department and University Extension, University of Western Australia, Publication No.8, p.143-166.
- Hausseux, M. (1990) Regional geology and geochemistry maps for the Meliadine River Project, Rankin Inlet, N.W.T.; Asamera Minerals Inc. unpublished maps 46, 47, 48 and 52.
- Helmstaedt, H.H. (1993) Natural diamond occurrences and tectonic setting of "primary" diamond deposits; *In Proceedings of a short course presented by the Prospectors and Developers Association of Canada*; March 27, 1993, Toronto, Ontario, p.3-72.

- McMartin, I. (2000) Till composition across the Meliadine trend, Rankin Inlet area, Kivalliq region, Nunavut; Geological Survey of Canada, Open File 3747.
- McMartin, I. And Henderson, P.J. (1999) A relative ice-flow chronology for the Keewatin Sector of the Laurentide Ice Sheet, Northwest Territories (Kivalliq Region, Nunavut); in Current Research, Geological Survey of Canada, Paper 1999-C, p. 129-138.
- Mitchell, R.H. (1986) Kimberlite: Mineralogy, Geochemistry and Petrology. Plenum Press, New York, 442 pages.
- Mitchell, R.H. (1989) Aspects of the petrology of kimberlites and lamproites: some definitions and distinctions; *In* Kimberlites and Related Rocks, Volume 1, Their Composition, Occurrence and Emplacement; Geological Society of Australia, Special Publication No.14, p.7-46.
- Mitchell, R.H. (1991) Kimberlites and lamproites: Primary sources of diamond. Geoscience Canada, Vol.18, p.1-16.
- Reinhardt, E.W. and Chandler, F.W. (1973) Gibson-MacQuoid Lake map area, District of Keewatin; in Report of Activities, Part A, Geological Survey of Canada, Paper 73-1A, p. 162-165.
- Scott Smith, B.H. (1995) Petrology and diamonds. Exploration and Mining Geology, Vol.4, No.2, p.127-140.
- Seller, M.H. (1999) Petrology of the Meliadine kimberlite dykes, District of Keewatin, Northwest Territories, Canada; M.Sc. Thesis, University of Alberta, Spring 1999, 227 p.
- Shilts, W.W., Cunningham, C.M., and Kaszycki, C.A. (1979) Keewatin Ice Sheet – Re-evaluation of the traditional concept of the Laurentide Ice Sheet; Geology, v. 7, p. 537-541.
- Skinner, E.M.W. (1989) Contrasting Group I and Group II kimberlite petrology: towards a genetic model for kimberlites. *In* J. Ross (ed.) Kimberlites and Related rocks, Vol.1, Their Composition, Occurrence, Origin and Emplacement, Proceedings of the Fourth Kimberlite Conference, Perth, 1986, Geological Society of Australia, Special Publication No.14, p.528-544.
- Tella, S. and Annesley, I.R. (1987) Precambrian geology of parts of the Chesterfield Inlet map area, District of Keewatin; in Current Research, Part A, Geological Survey of Canada, Paper 87-1A, p. 25-36.

- Tella, S., Annesley, I.R., Borradaile, G.J., and Henderson, J.R. (1986) Precambrian geology of parts of Tavani, Marble Island and Chesterfield Inlet map areas, District of Keewatin, N.W.T.; Geological Survey of Canada, Paper 86-13, 20 p.
- Tella, S., Schau, M., Armitage, A.E., and Loney, B.C. (1993) Precambrian geology and economic potential of the northeastern parts of the Gibson Lake (55N) map area, District of Keewatin, Northwest Territories; in Current Research, Part C, Geological Survey of Canada, Paper 93-1C.
- Tella, S., Schau, M., Armitage, A.E., Seemayer, B.E., and Lemkow, D. (1992) Precambrian geology and economic potential of the Meliadine Lake-Barbour Bay region, District of Keewatin, Northwest Territories; in Current Research, Part C; Geological Survey of Canada, Paper 92-1C, p. 1-11.