



Arsenic freshwater toxicity data for phytoplankton species used in development of arsenic water quality guideline. EC50 refers to concentration at which 50% of organisms exhibit an effect on growth. VSUE refers to “very severe unfavourable effect”.

Source: Lorax 2005

AS A MUTUAL PROTECTION TO OUR CLIENT, THE PUBLIC AND OURSELVES, ALL REPORTS AND DRAWINGS ARE SUBMITTED FOR THE CONFIDENTIAL INFORMATION OF OUR CLIENT FOR A SPECIFIC PROJECT AND AUTHORIZATION FOR USE AND/OR PUBLICATION OF DATA STATEMENTS CONCLUSIONS OR ABSTRACTS FROM OR REGARDING OUR REPORTS AND DRAWINGS IS RESERVED PENDING OUR WRITTEN APPROVAL.

SCALE:	N/A	DESIGNED:	HHH
DATE:	MARCH 2005	CHECKED:	HHH
DRAWN:	SLF	APPROVED:	HHH

CLIENT:



BGC

BGC Engineering Inc.
AN APPLIED EARTH SCIENCES COMPANY

Calgary, Alberta.

Phone: (403) 250-5185

PROJECT

ULU MINE-WASTE ROCK & ORE STORAGE PLAN

TITLE

ARSENIC FRESHWATER TOXICITY DATA

PROJECT No.

0385-002-02

FIGURE No.

13

REV.

0